

What people share about the COVID-19 outbreak on Twitter? An exploratory analysis

Dhivya Karmegam , Bagavandas Mapillairaju

To cite: Karmegam D, Mapillairaju B. What people share about the COVID-19 outbreak on Twitter? An exploratory analysis. *BMJ Health Care Inform* 2020;**27**:e100133. doi:10.1136/bmjhci-2020-100133

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjhci-2020-100133>).

Received 03 February 2020
Revised 02 November 2020
Accepted 05 November 2020

ABSTRACT

Background The recent outbreak of respiratory illness caused by COVID-19 in Wuhan, China, has received global attention as it has infected thousands of individuals there, and later it has also been reported from other countries internationally. This study aims at performing an exploratory study on Twitter to understand the information shared among the community regarding the COVID-19 outbreak.

Methods COVID-19 related tweets were collected from Twitter using keywords from 18 January to 25 January 2020. Top-ranking tweets were taken as samples and then categorised based on the content. Expressions or opinion tweets were analysed qualitatively to understand the mindset of the people regarding the outbreak. Theme wise reachability evaluation of the messages was also performed.

Results Based on the content of the tweets, five themes were evolved: (1) general information; (2) health information; (3) expressions; (4) humour and (5) others. 57.42% of messages are general information followed by expressive tweets (24.12%). Humorous messages were liked the most, whereas health information tweets were retweeted the maximum. Fear was the predominant emotion expressed in the messages.

Conclusion The results of the study would be useful to focus on the dissemination of the right information and effective communication on Twitter related to health and outbreak management.

INTRODUCTION

In recent days, COVID-19 has conquered the headlines in the news media and captured global attention as it infected more than thousands in China, and also it has been spread to other countries globally. COVID-19 (also called 2019-nCoV) causes severe respiratory illness, and it was first reported and detected in Wuhan City, China, in December 2019. As of 26 January 2020, COVID-19 infections were also reported from many other countries that include USA, Australia, Japan, France and so on.¹ Controlling and preventing the outbreak need continuous monitoring and international collaboration.² There is a need for quick access to information regarding the outbreak, and the public

should be aware of the risks and challenges involved in managing and preventing the COVID-19 infection.³ At the beginning of the outbreak, quick communication of information to the public is vital to prevent the fast-spreading virus.⁴ Communication regarding health during the outbreak will be effective only if it is done from the perspective of the people and also the content should be relevant to the targeted people.⁵ Understanding the perception of the people regarding the virus infection and outbreak by traditional data collection techniques has a lot of practical difficulties. It consumes a lot of time and money. When there is a crisis, people tend to use social media to seek information and express their views and feelings.^{6,7} In the case of COVID-19 outbreak, when physical distancing, isolation and travel restrictions are imposed, social media turns out to be the forum for public discussion more than usual.⁸ Messages shared by the people in social media like Twitter can be used to understand the expectations of the people and their mindset. In turn, this will help in planning, communicating and managing the crisis effectively based on the needs of the public.⁹ Social media has the potential to assess the awareness level and the state of mind of the public regarding any particular unexpected event.^{10,11} Previous studies also confirm that such analysis helps the public health stakeholders to know about information dissemination immediately during such an infectious disease outbreak.¹²⁻¹⁴ The earlier researches confirm that social media is one of the effective tools used to communicate information regarding preventive measures during the outbreak.¹⁵⁻¹⁷ The analysis of the content of the messages in social media provides an insight regarding the awareness and the needs of the public.^{18,19} There is a gap between the information spread over by the media and the concerns of the people. The purpose of this study is to understand the information and



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Centre for Statistics, SRM Institute of Science and Technology, Kattankulathur, Tamil Nadu, India

Correspondence to

Dhivya Karmegam;
dhivya.megam@gmail.com

content communicated on Twitter from 18 January to 25 January 2020 by the people globally about the COVID-19 outbreak. This helps the public health stakeholders to plan Twitter communication and information dissemination regarding the outbreak and associated information. To use social media to disseminate the right information regarding COVID-19 pandemic to reach a wider public, the initial and most important step is to understand the content of the messages posted and shared by the people. The objectives of this study are:

1. To categorise the Twitter messages related to COVID-19 infection based on their content into themes.
2. To explore the opinions and emotions of the people regarding the COVID-19 outbreak qualitatively on Twitter.

METHODS

Twitter messages were collected from Twitter search Application Programming Interface using hashtags related to the COVID-19 outbreak for 7 days from 18 January to 25 January 2020. Original tweets (excluding retweets) in English were collected using the following hashtags: *coronavirus*, *wuhan*, *coronaviruswuhan*, *coronaoutbreak*, *wuhanvirus*, *WuhanCoronavirus*, *WuhanPneumonia*, *ChinaCoronaVirus*, *ChinaWuHan*, *2019nCoV*, *WuhanOutbreak*, *VirusCorona*, *nCoV2019* and *nCoV*. The collection of tweets using these keywords was done using R programming, and it was 100 500 tweets from 61 216 unique users. Among the tweets collected, only 2409 tweets were geo-referenced. During the considered period, maximum numbers of confirmed cases of COVID-19 were reported from the countries that do not use Twitter or from non-native English-speaking countries.²⁰ These above-mentioned limitations restricted us to further analyse the tweets based on their geo-location.

Sample tweets were extracted from collected tweets by top-ranked sampling strategy.²¹ The favourite count (number of times the message was liked) and the retweet count (number of times the message was shared) of a particular tweet is the indicator of reachability of the tweets.²² The higher the favourite or the retweet count, the higher will be the reachability of the message in the public. In this study, the tweets with the favourite or the retweet count greater than 100 would be considered as top-ranked tweets, and they were taken as samples (n=1219).

In order to understand the details and information shared by the people about outbreak in Twitter, content analysis was performed. Based on the content of the messages, themes (what is discussed in the tweet) were identified and coded by the direct content analysis method.^{23 24} The first author framed a codebook of themes with reference to the earlier studies^{25–27} and a general look at the messages. Based on the codebook, both the authors coded 100 randomly selected tweets independently. Inconsistencies in coding between the two authors were discussed and resolved while finalising

the codebook of themes. With reference to the finalised codebook, both the authors (DK and BM) assigned the themes to the sample tweets independently. All the differences in coding were discussed by both the authors together until an agreement in final coding was met. Twitter messages that contained more than one theme were assigned as such. Then, the theme-wise reachability of the messages was evaluated based on the favourite and retweet ratio as indicators. The retweeting frequency and number of times the tweets were liked (favourite count) were used as the measure of interactivity and reachability of the tweet.²² The retweet and favourite rate were calculated as a ratio by dividing the total retweet or favourite count corresponding to a particular theme by the total number of messages on that theme.

To understand the mindset of the people regarding the COVID-19 outbreak and to explore the emotions expressed regarding the situation, tweets that stated expressions and opinions were examined qualitatively by discourse analysis.^{28 29} As Twitter messages had a limited number of words with hidden information, discourse analysis was done. Expressive and opinion tweets were read many times by both the authors, and the categories that evolved from the text were identified. The categories identified and the contexts behind the expressions were discussed qualitatively with example tweets.

RESULTS

Content analysis

Five themes were identified from the randomly selected 100 tweets. They were health information, general information and updates, expressions or opinions, sarcasm or humour, and others. Personal conversations and tweets that were not related to COVID-19 outbreak were categorised as a theme, others. Out of 1219 sample records, 98 (8.04%) tweets were coded under others category. The maximum number of messages contains general information and updates followed by expression or opinion tweets were many. Table 1 describes the themes with example tweets and the number of tweets in each category.

Online supplemental file 1 gives the status ID of the sample tweets, corresponding theme, favourite count and retweet count. Each tweet had a maximum of two themes.

Among the four major themes, sarcasm or humour had the lowest count but the highest favourite ratio (925). The sarcastic tweet's favorite ratio of 925 indicates that each sarcastic message was likely to be favourited 925 times on an average. Tweets from health information had the highest retweet ratio (577). The higher the retweet ratio, the higher will be the shares and reachability of that content to the Twitter users. Table 2 shows the theme-wise favourite count, retweet count, and favourite and retweet ratio.

Qualitative analysis

The tweets that stated opinions and expressions were analysed qualitatively to understand the mindset,

Table 1 Themes with description, example tweets and the number of tweets in each category

Themes	Description	Example tweets	Number of tweets (%)
Health information	Tweets that had information regarding transmission, symptoms, prevention and treatment for respiratory illness caused by COVID-19.	<i>#WuhanCoronavirus Symptoms of 2019-nCoV include: A fever Dry cough Shortness of breath Trouble breathing Dear all: frequent hand washing, covering coughs, and avoid touching your face https://t.co/jdHHu2O3Yn.</i>	138 (11.32)
General information and updates	Tweets that give updates on the count of confirmed cases and mortality in different countries. Tweets that share information regarding the steps carried out by the government in airports and hospitals to control and prevent outbreaks. Videos and images of the situation at hospitals in China.	<i>'China has locked down a city of 11 million to slow the spread of the new strain of coronavirus. Latest estimates: ▶ 17 dead ▶ 4k infected ▶ 22+ cities affected in 5–8 countries ▶ WHO will soon decide on a declaration of global public health emergency'.</i>	700 (57.42)
Expressions or opinions	Messages in which people expressed their opinions, emotions and clarifications regarding the outbreak.	<i>'What a nightmare. There is no way any city this size would be comfortable in lockdown'.</i>	294 (24.12)
Sarcasm or humour	Humorous or witty text, sarcastic images, videos and memes related to the COVID-19 outbreak.	<i>'a little late ... The coronavirus has already gone into "tourist mode", currently taking airline trips around the globe'.</i>	92 (7.55)

emotions and the context behind the emotional expression. The subcategories that emerged from opinion or expressive tweets were negative emotions, positive and thankful, suggestions and seeking information. As the results were presented qualitatively, the frequency of the tweets under each sub-category was specified using terms such as 'often', 'frequent' and 'some'.^{30 31} The opinions, emotions and the context of expressions were explained qualitatively under each subcategory with example tweets.

Negative emotions

People expressed negative emotions such as fear, anger, sadness and frustration towards the outbreak situation. Fear was frequently revealed in the messages. Fear was expressed by the people on knowing the statistics on confirmed cases and deaths, transmission routes and also on watching live videos of hospitals. Some example tweets that indicated fear were given below.

The Videos Of People Collapsing On Chinese Streets Due To The Coronavirus Are Scary As Shit.

This is very bad news, meaning airports checkpoints don't work since some #coronavirus patients are

asymptomatic and will continue spreading 2019-nCoV without even knowing. Nightmare scenario! SCARY! South China Morning Post reports that 41 people are DEAD from coronavirus in China with more than 1000 people are infected.

Followed by fear, sadness was sensed often in the tweets. Sadness was expressed concerning the outbreak circumstances, affected people and deaths. Few sample tweets that show sadness were presented.

i swear it was only last week it had breached no national boundaries. now it's breached 7? that seems like a pretty rapid acceleration, how worried are we supposed to be.

My heart breaks for the poor people dying of the Coronavirus. They're finding them all over.

it's pretty sad that today is Chinese New Year's Eve and it's time for family members to gather around. Maybe this doctor hasn't seen his family for a long time... I'm sure he is really desperated...

Table 2 Theme-wise favourite count, retweet count, and favourite and retweet ratio

Themes	Favourite count	Retweet count	Favourite ratio	Retweet ratio
Health information	89 585	79 578	649	577
General information and updates	382 474	258 619	546	369
Expressions or opinions	259 600	87 287	883	297
Sarcasm or humour	85 098	24 851	925	270

Frustration, anger and concerns were also exhibited in some of the messages along with other emotions in the context of outbreak settings. In some tweets, people stated that there was a delay in the actions taken by the government in controlling the spread. Some tweets also disclosed the lack of trust in the initiatives taken by the government and other authorised institutions to control the outbreak. Discontent regarding the rumours and misinformation spread in the social media was also voiced in some of the tweets. Some of the sample tweets that expressed anger and frustration were given.

this coronavirus shit is nuts.

there's a new fuckin disease that's already killing people and is confirmed to be in the United States.

LIES!!! Why are we allowing people from Wuhan in Toronto after the quarantine was announced? This is irresponsible!!

let's just spread a bunch of disinfo about the coronavirus who fucking cares anymore.

Positive and thankful

Positive opinions and supportive thoughts were shared in some Twitter messages towards the action taken by the officials and stakeholders while handling the crisis. People also expressed their gratitude to the public health professionals for their patience and their concern.

#CoronavirusOutbreak in #Wuhan goes viral and touches many; respects and tributes go to all of those doctors, nurses, and scientists.

What wasn't inevitable: screening on presentation to the ER, immediate isolation, calling @TOPublicHealth. The fact that this occurred shows the system is working. Kudos to staff at @Sunnybrook for your diligence.

Suggestions

In some messages, the public stated their personal opinions and suggestions regarding what could be done to control the virus spread and outbreak. One of the frequent suggestions mentioned in messages was closing the international borders thereby restricting people from other countries. Sample opinion tweets were given below.

Now it's spreading in the United States. Two words, TRAVEL BAN.

With Coronavirus Spreading, Canada Must Restrict Incoming Flights From China.

Seeking information

People also look for information and asks for clarification in relation to the outbreak. The public seeks information regarding the prevention and treatment, updates on affected cases and government initiatives. Twitter messages in which people asked questions regarding the outbreak were provided.

So when does the WHO declare coronavirus an emergency?

Anyone have a total confirmed coronavirus case number?

Is sushi safe to eat with this coronavirus shit going on?

DISCUSSION

The main aim of this study was to understand the content of the Twitter messages regarding the COVID-19 outbreak. From the results, it was clear that a maximum of general information and updates regarding the virus and outbreak was shared in social media. The next highest content in the Twitter messages was personal opinions or expressions shared by the people. This implies that people were aware of and also concerned about the spread of COVID-19 infections. Although the considered period in this study was the initial stage of the pandemic, we see a significant count of health information posted and shared. Health information messages posted would have provided the public, the knowledge regarding the symptoms and the prevention of COVID-19. Based on the reachability evaluation, even though sarcastic or humorous messages were liked the most, health information messages were likely to be retweeted and shared the maximum. This may be because people would have thought sharing health information was vital so that it reaches many Twitter users. However, regardless of the situation, people enjoyed the humorous tweets and memes posted.

Based on our qualitative analysis, we see that the public expressed mixed emotions and opinions (both positive and negative). Fear and sadness were predominantly prevailing among the people. The live videos of people suffering in hospitals, images of crowded hospitals, ambulances waiting with patients outside the hospital, incorrect personal opinions and information with more hyped numbers of the infected and mortality than the actual were widely shared in the Twitter messages. Some articles shared on Twitter presented the COVID-19 as a deadly and killer virus. These videos, images and personal opinions may be the reason for panic and unrest among the people. The information seeking behaviour of the public approves the responsiveness and concern of the public towards the crisis.

The evidence of misinformation in social media was recognised widely.^{32 33} Also the users did not share the link from the authorised website and sources directly on social media.³⁴ Hence, it is vital to check the credibility of the messages to handle the situation efficiently. Studies say that unverified content and rumours were shared mostly by individual accounts when compared with other sources like non-governmental organisations, media and healthcare accounts.^{33 35} As per the experts, one of the best approaches to avoid misinformation in social media is to increase social media messages from the authorised and official sources.³⁶ Healthcare professionals, researchers and famous personalities can be used

as a source to load social media with trustworthy information.³⁷ This will increase the verified information in social media, so that information and opinions of individuals may not dominate the social media content. If the information shared by the official sources satisfies the expectations and concerns of the people, then there will be no space for misinformation becoming viral.³⁶

Themes and expressions extracted from the tweets in this exploratory analysis give a basic understanding of the content, people's expectations and concerns. Based on this, outbreak managers can strategise communication by sharing messages that respond to public concerns and feelings especially fear. Clear, short, practical and shareable messages in social media can easily reach the audience. Other features of communication like multimedia and videos with illustrated comics may also help in spreading the information faster.^{38,39} Although the potential of social media analysis in crisis management and communication through the media has widely drawn the attention of the researches, it has never been integrated into practice for outbreak management.⁴⁰ The public health organisations and stakeholders need to understand the content, communicating features and trends in the social media and take advantage of those features to strategise the right method of information dissemination to the wider audience. An analysis of content and expressions were provided in this study and that can be used while planning communication of verified information dissemination.

This study analysed the data collected in 1-week duration. Also only a sample of tweets was considered for analysis. So, there is a possibility of limitations in the results arrived at using the limited data. One source (Twitter) of information of public views was examined in this study. Furthermore, the tweets only in English were considered. So there is a chance of losing the information from the messages posted in other social networking sites and other languages. However, this initial effort of content analysis of outbreak-related tweets gives a basic understanding of the situation and helps in planning the information dissemination on Twitter and other social media during the outbreak.

CONCLUSION

The study gives an overview of what has been shared on Twitter regarding the outbreak. Looking at the crisis from the point of view of the people and understanding their mindset is vital for the stakeholders in planning the response.^{6,41} COVID-19 related tweets were rich sources of information and opinions. As more information about the virus and the outbreak were posted and shared on Twitter, the same can be used by the public health professionals to disseminate the right information at the time of crisis.⁴² The results arrived at after examining the expression tweets give a basic understanding of people's needs and their state of mind towards the outbreak. Instead of letting the different opinions of the people dominating

the content of social media, health communications on Twitter can be strategised in such a way that it was replaced by scientific knowledge. Precise and verified information about virus transmission and prevention, communicated at the right time to the people, will improve their behavioural response and also help in reducing fear. This study analysed the tweets at a global level, whereas examining the tweets at the national level integrated with network analysis will give more detailed information for decision making.

Contributors Conceptualisation: BM and DK. Methodology: BM and DK. Formal analysis: DK and BM. Investigation: BM. Original draft preparation (report): DK. Review and editing (Report): BM and DK. Supervision: BM.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Dhivya Karmegam <http://orcid.org/0000-0003-3307-8704>

REFERENCES

- 1 CDC. Coronavirus disease 2019 (COVID-19) | centers for disease control and prevention, 2020. Available: <https://www.cdc.gov/coronavirus/2019-nCoV/summary.html> [Accessed 30 Jan 2020].
- 2 Liu S-L, Saif L. Emerging viruses without borders: the Wuhan coronavirus. *Viruses* 2020;12:130–7.
- 3 Wang C, Horby PW, Hayden FG, et al. A novel coronavirus outbreak of global health concern. *Lancet* 2020;395:10223:470–3.
- 4 Smith RD. Responding to global infectious disease outbreaks: lessons from SARS on the role of risk perception, communication and management. *Soc Sci Med* 2006;63:3113–23.
- 5 Vaughan E, Tinker T. Effective health risk communication about pandemic influenza for vulnerable populations. *Am J Public Health* 2009;99:S324–32.
- 6 Odium M, Yoon S. Health information needs and health seeking behavior during the 2014–2016 Ebola outbreak: a Twitter content analysis. *PLoS Curr* 2018;10. doi:10.1371/currents.outbreaks.fa814fb2bec36e29b718ab6af66124fa. [Epub ahead of print: 23 Mar 2018].
- 7 Gaspar R, Gorrão S, Seibt B, et al. Tweeting during food crises: a psychosocial analysis of threat coping expressions in Spain, during the 2011 European EHEC outbreak. *Int J Hum Comput Stud* 2014;72:239–54.
- 8 Chiolerio A. Covid-19: a digital epidemic. *BMJ* 2020;368:m764.
- 9 Lin X, Spence PR, Sellnow TL, et al. Crisis communication, learning and responding: best practices in social media. *Comput Human Behav* 2016;65:601–5.
- 10 Kass-Hout TA, Alhinnawi H. Social media in public health. *Br Med Bull* 2013;108:5–24.



- 11 Lachlan KA, Spence PR, Lin X. Expressions of risk awareness and concern through Twitter: on the utility of using the medium as an indication of audience needs. *Comput Human Behav* 2014;35:554–9.
- 12 Fu K-W, Liang H, Saroha N, et al. How people react to Zika virus outbreaks on Twitter? A computational content analysis. *Am J Infect Control* 2016;44:1700–2.
- 13 Vijaykumar S, Nowak G, Himelboim I, et al. Virtual Zika transmission after the first U.S. case: who said what and how it spread on Twitter. *Am J Infect Control* 2018;46:549–57.
- 14 Lazard AJ, Scheinfeld E, Bernhardt JM, et al. Detecting themes of public concern: a text mining analysis of the centers for disease control and prevention's Ebola live Twitter CHAT. *Am J Infect Control* 2015;43:1109–11.
- 15 Fung IC-H, Fu K-W, Ying Y, et al. Chinese social media reaction to the MERS-CoV and avian influenza A(H7N9) outbreaks. *Infect Dis Poverty* 2013;2:1–12.
- 16 Carter M. How Twitter may have helped Nigeria contain Ebola. *BMJ* 2014;349:g6946.
- 17 Oh S-H, Lee SY, Han C. The effects of social media use on preventive behaviors during infectious disease outbreaks: the mediating role of Self-relevant emotions and public risk perception. *Health Commun* 2020:1–10.
- 18 Lenoir P, Moulahi B, Azé J, et al. Raising awareness about cervical cancer using twitter: content analysis of the 2015 #SmearForSmear Campaign. *J Med Internet Res* 2017;19:e344.
- 19 Diddi P, Lundy LK. Organizational Twitter use: content analysis of Tweets during breast cancer awareness month. *J Health Commun* 2017;22:243–53.
- 20 WHO. Novel Coronavirus (2019-nCoV) situation report - 5 25 January 2020. World Heal. Organ, 2020. Available: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200125-sitrep-5-2019-ncov.pdf?sfvrsn=429b143d_8 [Accessed 30 Jan 2020].
- 21 Le GM, Radcliffe K, Lyles C, et al. Perceptions of cervical cancer prevention on Twitter uncovered by different sampling strategies. *PLoS One* 2019;14:1–14.
- 22 Park H, Reber BH, Chon M-G. Tweeting as health communication: health organizations' use of Twitter for health promotion and public engagement. *J Health Commun* 2016;21:188–98.
- 23 Hand RK, Kenne D, Wolfram TM, et al. Assessing the viability of social media for disseminating evidence-based nutrition practice guideline through content analysis of Twitter messages and health professional interviews: an observational study. *J Med Internet Res* 2016;18:e295–21.
- 24 Jiang X, Jiang W, Cai J, et al. Characterizing media content and effects of organ donation on a social media platform: content analysis. *J Med Internet Res* 2019;21:e13058.
- 25 Chew C, Eysenbach G. Pandemics in the age of Twitter: content analysis of Tweets during the 2009 H1N1 outbreak. *PLoS One* 2010;5:e14118–13.
- 26 Ahmed W, Bath PA, Scaffi L. Moral panic through the lens of Twitter: An analysis of infectious disease outbreaks. In: *ACM International Conference proceeding series*. Association for Computing Machinery, 2018: 217–21.
- 27 Sutton J, Vos SC, Olson MK, et al. Lung cancer messages on twitter: content analysis and evaluation. *J Am Coll Radiol* 2018;15:210–7.
- 28 Rachel Henzell M, Margaret Knight A, Morgaine KC, Henzell MR, Knight AM, et al. A qualitative analysis of orthodontic-related posts on Twitter. *Angle Orthod* 2014;84:203–7.
- 29 Marwick AE. *Ethnographic and qualitative research on Twitter*. Twitter and Society, 2014: 109–22.
- 30 Hays R, Daker-White G. The care.data consensus? A qualitative analysis of opinions expressed on Twitter health policies, systems and management in high-income countries. *BMC Public Health* 2015;15:1–13.
- 31 Bethea J, Murtagh B, Wallace SE. "I don't mind damaging my own body" A qualitative study of the factors that motivate smokers to quit. *BMC Public Health* 2015;15:1–9.
- 32 Sommariva S, Vamos C, Mantzarlis A, et al. Spreading the (fake) news: exploring health messages on social media and the implications for health professionals using a case study. *Am J Health Educ* 2018;49:246–55.
- 33 Kouzy R, Abi Jaoude J, Kraitem A, et al. Coronavirus goes viral: quantifying the COVID-19 misinformation epidemic on Twitter. *Cureus* 2020;12:e7255.
- 34 Bond CS, Ahmed OH. Can I help you? information sharing in online discussion forums by people living with a long-term condition. *J Innov Health Inform* 2016;23:620–6.
- 35 Wang Y, McKee M, Torbica A, et al. Systematic literature review on the spread of health-related misinformation on social media. *Soc Sci Med* 2019;240:112552.
- 36 Llewellyn S. Covid-19: how to be careful with trust and expertise on social media. *BMJ* 2020;368:m1160.
- 37 Tang L, Bie B, Park S-E, et al. Social media and outbreaks of emerging infectious diseases: a systematic review of literature. *Am J Infect Control* 2018;46:962–72.
- 38 Gray NA, Back AL. Covid-19 communication AIDS. *BMJ* 2020;369:m2255.
- 39 Yoon J, Hagen L, Andrews J, et al. On the use of multimedia in Twitter health communication: analysis of tweets regarding the Zika virus. *Inf Res* 2019.
- 40 Charles-Smith LE, Reynolds TL, Cameron MA, et al. Using social media for actionable disease surveillance and outbreak management: a systematic literature review. *PLoS One* 2015;10:e0139701.
- 41 Odium M, Yoon S, Broadwell P, et al. How Twitter can support the HIV/AIDS response to achieve the 2030 eradication goal: in-depth thematic analysis of World AIDS day Tweets. *JMIR Public Health Surveill* 2018;4:e10262.
- 42 Madanian S, Parry DT, Airehrour D, et al. mHealth and big-data integration: promises for healthcare system in India. *BMJ Health Care Inform* 2019;26:e100071.