Technology report

Design and implementation of an affordable, public sector electronic medical record in rural Nepal

Anant Raut* and Bikash Gauchan

Possible, Bayalpata Hospital, Achham, Nepal

Chase Yarbrough* and Dan Schwarz

Possible, Bayalpata Hospital, Achham, Nepal, Brigham and Women's Hospital, Department of Medicine, Division of Global Health Equity, Boston, USA, and Boston Children's Hospital, Department of Medicine, Division of General Pediatrics, Boston, USA

David Citrin

Possible, Bayalpata Hospital, Achham, Nepal, Department of Anthropology, University of Washington, Seattle, USA, Department of Global Health, University of Washington, Seattle, USA, and Henry M. Jackson School of International Studies, University of Washington, Seattle, USA

Vivek Singh

ThoughtWorks India, Bengaluru, India

Varun Verma

Mount Sinai Hospital, Department of Internal Medicine, Division of Hospital Medicine, New York, USA, and Department of Internal Medicine, Division of Hospital Medicine, Icahn School of Medicine at Mount Sinai, New York, USA

Jessica Hawley

UCLA Medical Center, Santa Monica, USA, and Department of Medicine, Division of Hospitalist Services, David Geffen School of Medicine at UCLA, Los Angeles, USA

Alex Harsha Bangura

Contra Costa Regional Medical Center, Martinez, USA

Biplav Shrestha

Patan Hospital, Department of Pediatric Research, Lalitpur, Nepal

Ryan Schwarz

Possible, Bayalpata Hospital, Achham, Nepal, Brigham and Women's Hospital, Department of Medicine, Division of Global Health Equity, Boston, USA, Department of Medicine, Harvard Medical School, Boston, USA, and Massachusetts General Hospital, Department of Medicine, Division of General Internal Medicine, Boston, USA

Mukesh Adhikari

Ministry of Health, Department of Health Services, District Health Office, Achham, Nepal

Duncan Maru

Possible, Bayalpata Hospital, Achham, Nepal, Brigham and Women's Hospital, Department of Medicine, Division of Global Health Equity, Boston, USA, Boston Children's Hospital, Department of Medicine, Division of General Pediatrics, Boston, USA, and Department of Global Health and Social Medicine, Harvard Medical School, Boston, USA, and Department of Medicine, Harvard Medical School, Boston, USA

ABSTRACT

Introduction Globally, electronic medical records are central to the infrastructure of modern healthcare systems. Yet, the vast majority of electronic medical records have been designed for resource-rich environments and are not feasible in settings of poverty. Here, we describe the design and implementation of an electronic medical record at a public sector district hospital in rural Nepal and its subsequent expansion to an additional public sector facility.

Development The electronic medical record was designed to solve for the following elements of public sector healthcare delivery: 1) integration of the systems

*Anant Raut and Chase Yarbrough are co-first authors.

Cite this article: Raut A, Yarbrough C, Singh V, Gauchan B, Citrin D, Verma V, Hawley J, Schwarz D, Harsha Bangura A, Shrestha B, Schwarz R, Adhikari M, Maru D. Design and implementation of an affordable, public sector electronic medical record in rural Nepal. *J Innov Health Inform.* 2017;24(2):186–195.

<http://dx.doi.org/10.14236/jhi.v24i2.862>

Copyright © 2017 The Author(s). Published by BCS, The Chartered Institute for IT under Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

Author address for correspondence

Duncan Maru
Division of Global Health Equity
Boston, USA
Email: duncan@possiblehealth.org

Accepted March 2017

across inpatient, surgical, outpatient, emergency, laboratory, radiology and pharmacy sites of care; 2) effective data extraction for impact evaluation and government regulation; 3) optimization for longitudinal care provision and patient tracking and 4) effectiveness for quality improvement initiatives.

Application For these purposes, we adapted Bahmni, a product built with open-source components for patient tracking, clinical protocols, pharmacy, laboratory, imaging, financial management and supply logistics. In close partnership with government officials, we deployed the system in February of 2015, added on additional functionality and iteratively improved the system over the following year. This experience enabled us then to deploy the system at an additional district-level hospital in a different part of the country in under four weeks. We discuss the implementation challenges and the strategies we pursued to build an electronic medical record for the public sector in rural Nepal.

Discussion Over the course of 18 months, we were able to develop, deploy and iterate upon the electronic medical record, and then deploy the refined product at an additional facility within only four weeks. Our experience suggests the feasibility of an integrated electronic medical record for public sector care delivery even in settings of rural poverty.

Keywords: electronic medical records, health systems strengthening, implementation research, global health, Nepal, open source technologies

INTRODUCTION

The digitization of healthcare delivery systems is a pressing global need even in settings of poverty.^{1,2} The opportunity costs of continuing with paper systems are significant. The absence of electronic medical records (EMRs) can hamstring already challenging efforts to transform human resource management, financial accountability, healthcare systems performance evaluation, public health surveillance and longitudinal care delivery systems.^{3,4} Paper-based systems are human resource and time intensive and are often plagued by inaccurate reporting processes leading to out-of-date and irrelevant data.^{3–5} An effective EMR is an essential component of a robust and efficient modern healthcare system.^{3,6,7}

Despite the strong evidence that adoption of health information technology leads to improved care,^{8,9} only recently have wealthy countries started to see widespread adoption.¹⁰ Low- and middle-income countries face additional challenges such as a lack of engineering and other technical expertise, limited funding, poor information technology infrastructure and unreliable power.^{2,11} Indeed, many hospital information systems that have been successfully implemented in resource-poor countries tend to be focused on specific clinical diseases rather than care integration,¹² and there are only limited descriptions of such integrated systems.¹³

Even the most sophisticated and well-funded systems have achieved greater success in data recording, research, and program evaluation than in creating a tool that physicians can refer to during a clinical encounter.^{2,14} Although there have been efforts to address these challenges,^{14–17} successful implementation of integrated EMR systems, particularly

those that are actively used by healthcare providers during clinical encounters, has been an elusive goal.

DEVELOPMENT

Possible, a non-profit healthcare organization, started in June 2014 to customize and deploy an EMR system appropriate for the Nepali public sector healthcare system. *Possible* works on a public–private partnership model with the Ministry of Health in rural Achham district in the far-western development region. There, *Possible* independently manages the government-owned, district-level Bayalpata Hospital and implements community healthcare delivery programs. Bayalpata Hospital has 25 beds and sees approximately 60,000 outpatients and 2,000 inpatients per year. Patients are seen by physicians and health assistants, a mid-level non-physician provider role common throughout Nepal.

We originally identified the following key needs as central to developing an EMR: to be simple for physicians and mid-level providers, to improve government reporting, to decrease wasteful expenditures on unnecessary medications, to use the EMR for healthcare services research, and to advance quality improvement initiatives. After some review and deliberations in the first months of 2014, *Possible* selected *Thought Works*' (Chicago, USA) Bahmni system as its design characteristics were suited to the on-the-ground realities of Achham (Box 1).

Bahmni (<http://www.Bahmni.org>) is an open source framework built as a javascript application on top of the OpenMRS (<http://www.openmrs.org>) data model and application program interface. Bahmni uses OpenERP (<https://www.odoo.com/>) for billing and inventory management and OpenELIS (<http://www.openelis.org/>) for laboratory management.

Box 1 Design characteristics of the EMR system

Design Characteristics	
Longitudinal care	The product needs to be designed to enhance the goal of improved longitudinal care, including both patient tracking and performance quality monitoring. Simple disease management protocols and data tracking should be modularized and accessible within the system.
Cross-Site Integration	The product should be integrated from the hospital, where human resource, electricity, and internet capacity are all substantial, to the primary clinics which are severely resource-constrained, to patients and community health workers, who would interface with the product on mobile devices.
Scalability	In everything we do, we have to design for scale. This is particularly true because there is a strong push for us to expand the idea of public-private partnerships where the government is a payer and regulator and Possible is the direct implementer. If the technology can't ultimately be scaled at a national level, then it is not worth the effort.
Reliability	Patients will suffer needless delays in care if the system is not exquisitely reliable in the setting in which the product is deployed.
Offline capability	The product needs to be online/offline capable, or at least intranet-capable, since, while Possible can get computers electricity 95% of the time, the internet (despite serious investments) is only 85% at best.

Thought Works has been developing Bahmni over the last four years for deployments in India, Uganda, Haiti and other countries. We provide a diagram of the system architecture in Figure 1.

A primary innovation of the Bahmni system is that it provides a stripped-down user-friendly interface for the robust and widely used OpenMRS architecture. In a resource-limited setting such as rural Nepal, most providers have had limited prior exposure to computers or EMR systems. A well-tuned user-experience without non-essential features is thus of critical importance for providers to be able to use the EMR at the point of care without feeling overwhelmed. Workflow efficiency in the outpatient department is paramount, since Bayalpata Hospital can care for over 600 outpatients per day during peak seasons (Figure 2). *Possible* uses the EMR in the inpatient department for bed management, drug administration, laboratory orders and results review, progress notes documentation, and discharge planning and documentation. In the operating theaters, *Possible* uses the EMR to document operative notes.

APPLICATION

Bayalpata hospital EMR implementation

EMR implementation necessarily took place in a phased manner (Box 2). In preparation for the initial rollout of Bahmni at Bayalpata Hospital, a team from *Thought Works* came to the hospital and together with a staff trainer, trained the doctors, mid-level providers, nurses, and registration staff over a period of two weeks in July of 2014. Over the subsequent six months, we undertook several off-site and on-site design iterations and planning sessions. On 10 February 2015, Bahmni went live at the hospital with the patient registration

Figure 1 Bahmni architecture

*PACS = Picture Archiving and Communication System; USG/ECG = Ultrasound Sonography Test/ Electrocardiography; DICOM = Digital Imaging and Communications in Medicine; ICD-10 = International Classification of Diseases, version 10

Figure 2 Patient workflow at Bayalpata Hospital

module. The EMR was deployed to the laboratory and outpatient departments over the following week. This phased roll-out allowed the deployment team to adequately support each department during the initial stage. Initially, providers were only required to enter diagnoses for patients, but over the course of time, these providers became able to review medical history and lab results, enter vitals, fill out notes, order lab tests, and prescribe medications from within the EMR.

Providers access the EMR on touchscreen Chromebooks made available at all points of care. Chromebooks were chosen because they are low maintenance, inexpensive (below USD \$300), and ideal for browser-based applications like Bahmni. Clinical notes have been designed to take advantage

of the touchscreen functionality through use of buttons where possible for quick data entry. Unlike with electronic tablets, a physical keyboard is available when typing is required (Figures 2a, 2b and 3).

EMR response and challenges

We conducted individual focus group discussions with three different participant groups – doctors, mid-level providers and nurses – three months after the deployment of the first phase. Five doctors, thirteen mid-level providers, and twelve nurses participated in the focus groups. We held a discussion with each group that began with directed questions and ended with open discussion. Participants also

Box 2 EMR implementation timeline with 3 release phases.

PHASES OF IMPLEMENTATION

*IPD = Inpatient Department; IMAM = Integrated Management of Acute Malnutrition; USG = Ultrasound Sonography Test; CHD = Community Health Department; DHIS2 = District Health Information Software, version 2

Figure 2a Example patient dashboard

completed an anonymous survey with questions scored on a scale from one to ten, as well an area for free-text comments (Table 1).

Overall, the respondents described the EMR as a useful tool that improved the hospital and had the potential of making their work easier in the future. The focus group discussions and free-text responses, however, elicited several challenges. The doctors expressed concerns that introducing a computer into the room distracted from building the patient-provider relationship. The mid-level providers and doctors raised the issue that the EMR led to extra documentation time; when the power went out, they were forced to document encounters on paper and again later in the EMR. The nursing focus group revealed that there was some confusion about documentation responsibilities and the responsibilities of nurses in documenting. Many participants expressed challenges that parallel those of others working with EMRs in wealthier countries.^{18,19}

An anticipated challenge, one that was persistently difficult to mitigate, was regulating the server room's temperature. The servers require maintenance of temperatures below 23°C, yet during the summer months, ambient temperatures frequently reach 35°C. This required installation of air conditioner-heater systems that have significant power needs. Like many district hospital systems, at the time of predeployment, the hospital was already operating beyond

its electric power capacity. This issue caused occasional EMR down times, which challenged clinicians' faith and patience with the system. Bayalpata Hospital addressed these issues with ongoing investments in grid improvements, solar power, generators, battery backups, and more effective load balancing. It was difficult, however, to keep up with the demands on reliable electricity throughout the roll-out period.

Human resource management and training have also been core challenges. Most of the end-users had only minimal exposure to computers and no prior experience with EMRs. Initially, users voiced some discontent about an increased time investment per patient. Becoming adept at the EMR system and to the computers themselves took focus and time away from the patient. *Possible* designated a staff physician as the lead clinical point person to help coordinate trainings, make announcements on system changes, and gather feedback and suggestions from the care delivery team on how to improve the functionality and interface. The use of a champion from within the clinical staff helped with encouraging, processing, and utilizing feedback from front-line providers. Over several months, the clinicians developed significant improvement in their ability to efficiently navigate Bahmni.

Following this model, *Possible* selected focal personnel from different departments within the hospital (e.g. nursing,

▼ Acute Abdominal Complaint

Chief Complaint

Duration

▼ Symptoms

Pain characteristic

Location

Mid epigastric

RUQ

LUQ

RLQ

LLQ

Radiation

Nausea/Vomiting

Yes

No

Diarrhoea

Yes

No

Blood in vomit or stool

Yes

No

Other Symptoms

Water brash/burping

Difficulty swallowing

Fever

Loss of appetite

Weight Loss

Other significant history

LMP

Allergies

► Vitals

+ Add Section

Physical Examination Details

▼ Problem/Plan

Problem

Plan

Anti-infective

H2 blocker

PPI

Ultrasound

Stool

ER

Follow up in

SOS

3 days

5 days

1 week

2 weeks

1 month

>1 month

Figure 2b Example EMR clinical note

surgery, mental health and pharmacy) to form an EMR-focused clinical informatics team at Bayalpata Hospital. This team meets quarterly to discuss ways to further optimize the EMR. Additionally, Bahmni and the OpenMRS development

communities have created a system and culture responsive to user feedback, and as a result, *Thought Works* continues to enhance the user interface to make Bahmni as intuitive and efficient as possible. In order to take full advantage of the

A photograph showing a male clinician with dark hair and glasses, wearing a blue shirt, sitting at a desk and using a laptop. The desk is cluttered with various medical supplies, including boxes of FaceMask, a box of 50% alcohol hand sanitizer, and other medical equipment. The laptop screen displays a web-based interface, likely the EMR system.

Figure 3 Clinician at Bayalpata Hospital using a Chromebook

Table 1 EHR Focus Group Survey Responses. Doctors, health assistants and nurses were surveyed on their opinion of the EMR. There were eight questions scored on a scale of 1 to 10

Question	Score (average)
On a scale of 1 to 10, how useful is the EHR for you right now?	7.9
On a scale of 1 to 10, how useful could the EHR be for you in the future?	9.0
I feel competent using the EHR	7.9
I like using the EHR	8.9
The EHR makes my work more difficult	3.2
The EHR makes my work easier	8.4
The EHR will provide useful data that will improve patient care	9.1
The EHR is good for Bayalpata Hospital	9.4

Journal of Innovation in Health Informatics Vol 24, No 2 (2017)

BMJ Health Care Inform: first published as 10.14236/jhi.v24i2.862 on 1 April 2017. Downloaded from <http://informatics.bmj.com/> on April 26, 2023 by guest. Protected by copyright.

EMR system, work still remains to educate providers, habituated in paper-based systems, to write good notes and use the data for better clinical decision-making.

EMR refinement

Constant iteration is necessary in these early implementation phases to optimize the system for usability and utility, ensuring that the EMR is providing increasing value to patient care, and meaningful data for program management, evaluation, and quality improvement.

Our focus thus far has been on setting up a simple yet robust EMR system and building end-user capacity to take advantage of the system. Our efforts at monitoring and evaluation have centred on increasing EMR use and improving data quality. To this effect, we measure indicators such as 'percent of patients who were discharged without a discharge note in the EMR' and 'percent of outpatients without a treatment plan documented' (Figure 4). We also review manually entered non-coded diagnoses and perform daily data quality sweeps. These efforts continue in earnest as we are increasingly using the system to monitor indicators directly related to patient care and patient experience, e.g. we are currently refining our approach to measuring outpatient cycle time and care-to-wait-time ratios using time-stamped data pulled from the EMR in a quality improvement dashboard (Figure 5). Using data generated from the EMR, we have rolled out indicators that monitor chronic disease-specific quality metrics, for example blood pressure, glycosylated hemoglobin (HbA1c) and CD4 targets.

Improvement in stock management at both the supply chain and facility levels are significant advantages of the EMR. The digital platform allows us to monitor the requisition and movement of medicines and other consumables, and flexible reporting systems enable quality improvement efforts (Figure 6). For example, we measure stock out and maintenance rates as well as the percentage of essential medicines received from the government. At Bayalpata Hospital, drug

Figure 4 Patient indicators generated from EMR data

orders by providers are sent directly to the pharmacy for dispensing, saving patients time, and orders are then automatically deducted from the hospital inventory.

Replication and scale

A major rationale for the OpenMRS-based Bahmni systems was that it is affordable (Table 2) and designed for the types of providers found among public healthcare institutions in South Asia. From the beginning, *Possible* engaged government partners at the local and national levels. Indeed, using an OpenMRS-based system like Bahmni was itself predicated on the Ministry of Health's commitment to OpenMRS as the back-end program for its healthcare

Figure 5 Quality improvement dashboard

Figure 6 Stock management pareto analysis

system. This engagement requires constant dialogue, as there are frequent shifts at the leadership levels within the Ministry. Policy priorities themselves have been dynamic, particularly after the devastating earthquakes on 25 April 2015 and 12 May 2015.

The national dialogue surrounding EMR standards and possibilities has been influenced by successful implementation of a full-service EMR in a rural district hospital, leading our government partners to wonder whether a similar system might be employed more broadly across government facilities. Subsequently, when Possible initiated a new public-private partnership at a district hospital in Dolakha, a district heavily affected by the earthquakes, Possible quickly and successfully deployed the EMR within four weeks.

The IT infrastructure required to host an open source EMR system like Bahmni requires initial investment in materials and expertise. In our experience, however, ongoing costs are much lower due to the availability of remote support, the lack of licensing fees since we have used open-source software, a substantial OpenMRS design and implementation community on the Web, and regional experts for maintenance of the system. These resources are available within Nepal, even in rural areas. Indeed, partly as a result of deployments in Achham and Dolakha, the government has made commitments to expand an OpenMRS-based EMR throughout the country, using private sector technology companies to deploy and maintain the systems.

Table 2 EHR implementation costs

#	Activity	Duration	Estimated cost (in USD)
1	Development, customization, rollout	11 months	\$160,000
2	Travel (analysis and rollout)		\$8,200
3	Post Production Support	1 year	\$10,000
4	Servers and networking		\$32,000
5	Chromebooks (29@ \$300)		\$8,700
	TOTAL (Estimated)		\$218,900

DISCUSSION

We have successfully implemented an affordable integrated EMR system at two public sector rural district hospitals in Nepal. Our initial deployments have proved acceptable to clinicians, technologically feasible, and compatible with the national healthcare information system. We are now refining and expanding the EMR and making preparations for scale. Eventually, the EMR will provide both patient-level and population-level outcomes for quality improvement, health systems research, and public health surveillance systems.

Acknowledgments

We wish to express our appreciation to the Government of Nepal, Ministry of Health, for their continued efforts to improve access to healthcare in rural Nepal and for their advocacy around the use of EMRs in the public sector health-care system. We also wish to express our deepest appreciation to our partners at the Achham District Health Office and Public Health Office and all of the allied healthcare workers serving Achham and Dolakha Districts. These include the Female Community Health Volunteers, (Sub) Health Post and Primary Care Center staff members, and District Health Office staff members. We also wish to thank Lena Wong for her assistance in preparing the figures.

Conflict of interest statement

AR, BG and DC are currently employed by a nonprofit healthcare company (Possible) that delivers free healthcare in rural Nepal using funds from the Government of Nepal and other public, philanthropic, and private foundation sources. CY, DS, RS and DM all work in partnership with Possible and VV, JH and AHB worked in partnership with Possible at the time of electronic health record implementation. CY, DS and DM have been employed at two academic medical centres (Brigham and Women’s Hospital and Boston Children’s Hospital) that receive public sector research funding as well as revenue through private sector fee-for-service medical transactions and private foundation grants. DC is a faculty member

and employed part time at a public university (University of Washington). VV is employed at an academic medical centre (Mount Sinai Hospital) that receives public sector research funding, as well as revenue through private sector fee-for-service medical transactions and private foundation grants. VV is also a faculty member at a private university (Icahn School of Medicine at Mount Sinai). JH is employed at an academic medical centre (UCLA Medical Center) that receives public sector research funding as well as revenue through private sector fee-for-service medical transactions and private foundation grants. JH is also faculty at a public university (UCLA). AHB is a medical resident at a public hospital (Contra Costa Regional Medical Center), which receives revenue through fee-for-service medical transactions and other grants. BS was employed by Possible at the time of electronic medical record implementation and is currently employed by an academic medical centre (Patan Hospital) which receives revenue through fee-for-service medical transactions and other public, philanthropic, and private foundation sources. RS is employed at two academic medical centers (Brigham and Women's Hospital and Massachusetts General Hospital) that receive public sector research funding, as well as revenue through private sector fee-for-service medical transactions and private foundation grants. MA was employed by the Government of Nepal. DM is a non-voting member on Possible's board of directors, but receives no compensation. RS and DM are also faculty members at a private university (Harvard Medical School). All authors have read and understood the Journal of Innovation in Health Informatics' Ethics and Publication Malpractice Statement, and declare that we

have no competing financial interests. The authors do, however, believe strongly that healthcare is a public good, not a private commodity.

Funding statement

Research reported in this publication was supported by the Office of the Director, National Institutes of Health of the National Institutes of Health, under Award DP5OD019894. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The Eunice Kennedy Shriver National Institute Of Child Health & Human Development and the National Institute of Dental & Craniofacial Research provided support for this award. Research reported in this publication was also supported by the Charles H. Hood Foundation. The funders had no role in the program design, program implementation, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions

Designed the EMR system: AR, CY, VS, DC and DM

Implemented the EMR system: all authors

Conducted focus group discussions: AR, DC and CY

Analyzed data and evaluated the EMR system implementation: AR, CY, DC and DM

Contributed to writing the manuscript: AR, CY, DC and DM

Reviewed manuscript and critically revised content: all authors

ICMJE criteria for authorship read and met: all authors

Agree with manuscript results and conclusions: all authors

REFERENCES

1. Webster PC. The rise of open-source electronic medical records. *Lancet* 2011;377(9778):1641–2. PubMed PMID: 21591284. Epub 2011/05/19. eng. [https://doi.org/10.1016/S0140-6736\(11\)60659-4](https://doi.org/10.1016/S0140-6736(11)60659-4).
2. Williams F and Boren SA. The role of the electronic medical record (EMR) in care delivery development in developing countries: a systematic review. *Informatics in Primary Care* 2008;16(2):139–45. PubMed PMID: 18713530. Epub 2008/08/21. eng. <https://doi.org/10.14236/jhi.v16i2.685>.
3. Shekelle PG, Morton SC and Keeler EB. Costs and benefits of health information technology. *Evidence Report/Technology Assessment* 2006;132(132):1–71. PubMed PMID: 17627328. Epub 2007/07/14. eng.
4. Uslu AM and Stausberg J. Value of the electronic patient record: an analysis of the literature. *Journal of Biomedical Informatics* 2008;41(4):675–82. PubMed PMID: 18359277. Epub 2008/03/25. eng. <https://doi.org/10.1016/j.jbi.2008.02.001>. PMID:18359277.
5. DesRoches CM, Campbell EG, Rao SR, Donelan K, Ferris TG, Jha A et al. Electronic medical records in ambulatory care – a national survey of physicians. *The New England Journal of Medicine* 2008;359(1):50–60. PubMed PMID: 18565855. Epub 2008/06/21. eng. <https://doi.org/10.1056/NEJMs0802005>. PMID:18565855.
6. Whittaker M, Hodge N, Mares RE and Rodney A. Preparing for the data revolution: identifying minimum health information competencies among the health workforce. *Human Resources for Health* 2015;13:17. PubMed PMID: 25889677. Pubmed Central PMCID: PMC4391110. Epub 2015/04/19. eng. <https://doi.org/10.1186/s12960-015-0002-x>. PMID:25889677. PMCID:PMC4391110.
7. Chaudhry B, Wang J, Wu S, Maglione M, Mojica W, Roth E et al. Systematic review: impact of health information technology on quality, efficiency, and costs of medical care. *Annals of Internal Medicine* 2006;144(10):742–52. PubMed PMID: 16702590. Epub 2006/05/17. eng. <https://doi.org/10.7326/0003-4819-144-10-200605160-00125>. PMID:16702590.
8. Bates DW, Leape LL, Cullen DJ, Laird N, Petersen LA, Teich JM et al. Effect of computerized physician order entry and a team intervention on prevention of serious medication errors. *Journal of the American Medical Association* 1998;280(15):1311–6. PubMed PMID: 9794308. Epub 1998/10/30. eng. <https://doi.org/10.1001/jama.280.15.1311>. PMID:9794308.
9. Kaushal R, Shojania KG and Bates DW. Effects of computerized physician order entry and clinical decision support systems on medication safety: a systematic review. *Archives of*

- Internal Medicine* 2003;163(12):1409–16. PubMed PMID: 12824090. Epub 2003/06/26. eng. <https://doi.org/10.1001/archinte.163.12.1409>. PMID:12824090.
10. Adler-Milstein J, DesRoches CM, Furukawa MF, Worzala C, Charles D, Kralovec P et al. More than half of US hospitals have at least a basic EMR, but stage 2 criteria remain challenging for most. *Health Affairs (Project Hope)* 2014;33(9):1664–71. PubMed PMID: 25104826. Epub 2014/08/12. eng. <https://doi.org/10.1377/hlthaff.2014.0453>. PMID:25104826.
 11. Luna D, Almerares A, Mayan JC, Gonzalez Bernaldo de Quiros F and Otero C. Health informatics in developing countries: going beyond pilot practices to sustainable implementations: a review of the current challenges. *Healthcare Informatics Research* 2014;20(1):3–10. PubMed PMID: 24627813. Pubmed Central PMCID: PMC3950262. Epub 2014/03/15. eng. <https://doi.org/10.4258/hir.2014.20.1.3>. PMID:24627813; PMCID:PMC3950262.
 12. Tomasi E, Facchini LA and Maia MF. Health information technology in primary health care in developing countries: a literature review. *Bulletin of the World Health Organization* 2004;82(11):867–74. PubMed PMID: 15640923. Pubmed Central PMCID: PMC2623061. Epub 2005/01/11. eng. PubMed PMID: 15640923. PMID:15640923; PMCID:PMC2623061.
 13. Fraser HS, Biondich P, Moodley D, Choi S, Mamlin BW and Szolovits P. Implementing electronic medical record systems in developing countries. *Informatics in Primary Care* 2005;13(2):83–95. PubMed PMID: 15992493. Epub 2005/07/05. eng. <https://doi.org/10.14236/jhi.v13i2.585>.
 14. Scholl J, Syed-Abdul S and Ahmed LA. A case study of an EMR system at a large hospital in India: challenges and strategies for successful adoption. *Journal of Biomedical Informatics* 2011;44(6):958–67. PubMed PMID: 21846508. Epub 2011/08/19. eng. <https://doi.org/10.1016/j.jbi.2011.07.008>. PMID:21846508.
 15. Waters E, Rafter J, Douglas GP, Bwanali M, Jazayeri D and Fraser HS. Experience implementing a point-of-care electronic medical record system for primary care in Malawi. *Studies in Health Technology and Informatics* 2010;160(Pt 1): 96–100. PubMed PMID: 20841657. Epub 2010/09/16. eng. PMID:20841657.
 16. Douglas GG, Gadabu OJ, Joukes S, Mumba, S, McKay MV, Ben-Smith, et al. Using touchscreen electronic medical record systems to support and monitor national scale-up of antiretroviral therapy in Malawi. *PLoS Medicine* 2010;7(8):e1000319.
 17. Anokwa Y, Allen C, Yarborough C, and Frasier H, Ed. Building a Better Clinician Experience in OpenMRS. Global Partners in Public Health Informatics, Washington, DC: Anokwa, 2008.
 18. Margalit RS, Roter D, Dunevant MA, Larson S and Reis S. Electronic medical record use and physician-patient communication: an observational study of Israeli primary care encounters. *Patient Education and Counseling* 2006;61(1):134–41. PubMed PMID: 16533682. Epub 2006/03/15. eng. <https://doi.org/10.1016/j.pec.2005.03.004>. PMID:16533682.
 19. Poissant L, Pereira J, Tamblyn R and Kawasumi Y. The impact of electronic medical records on time efficiency of physicians and nurses: a systematic review. *Journal of the American Medical Informatics Association* 2005;12(5):505–16. PubMed PMID: 15905487. Pubmed Central PMCID: PMC1205599. Epub 2005/05/21. eng. <https://doi.org/10.1197/jamia.M1700>. PMID:15905487; PMCID:PMC1205599.