Atrial fibrillation dashboard evaluation using the think aloud protocol

Simon de Lusignan,^{1,2} Harshana Liyanage ,¹ Julian Sherlock,¹ Filipa Ferreira,¹ Neil Munro,¹ Michael Feher,¹ Richard Hobbs¹

To cite: de Lusignan S, Liyanage H, Sherlock J, et al. Atrial fibrillation dashboard evaluation using the think aloud protocol. *BMJ Health Care Inform* 2020;**27**:e100191. doi:10.1136/bmjhci-2020-100191

Received 11 June 2020
Revised 19 August 2020
Accepted 24 August 2020

ABSTRACT

Background Atrial fibrillation (AF) is a common cardiac arrhythmia which is a major risk factor for stroke, transient ischaemic attacks and increased mortality. Primary care management of AF can significantly reduce these risks. We carried out an evaluation to assess the usability of an AF dashboard developed to improve data quality and the quality of care.

Method We developed an online dashboard about the quality of AF management for general practices of the Oxford Royal College of General Practitioners Research and Surveillance Centre network. The dashboard displays (1) case ascertainment, (2) a calculation of stroke and haemorrhage risk to assess whether the benefits of anticoagulants outweigh their risk, (3) prescriptions of different types of anticoagulant and (4) if prescribed anticoagulant is at the correct dose. We conducted the think aloud evaluation, involving 24 dashboard users to improve its usability.

Results Analysis of 24 transcripts received produced 120 individual feedback items (ie, verbalised tasks) that were mapped across five usability problem classes. We enhanced the dashboard based on evaluation feedback to encourage adoption by general practices participating in the sentinel network.

Conclusions The think aloud evaluation provided useful insights into important usability issues that require further development. Our enhanced AF dashboard was acceptable to clinicians and its impact on data quality and care should be assessed in a formal study.

Summary box

What is already known?

- ▶ Atrial fibrillation (AF) is a major risk factor for stroke, transient ischaemic attacks; Primary care management of AF can significantly reduce these risks.
- ▶ Clinical dashboards assist clinicians, and managers to monitor and improve the quality of care of atrial fibrillation.

What does this paper add?

- ▶ We demonstrated the use of the think aloud protocol for evaluating the usability of a dashboard used in a primary care setting.
- ▶ AF management choices and quality (prescribing) sections were found to be the most useful indicators for clinical practice.

the possibility of tools being successfully integrated into routine clinical workflows, providing greater efficiency, and ultimately in quality improvement.^{7 8}

Atrial fibrillation (AF) is one of the most common and important heart arrhythmias; if undetected and left untreated, it can result in stroke and increased mortality. Early recognition of AF in practice can lead to early intervention with managing the risks of these complications. Current guidelines on the management of AF by National Institute for Health and Care Excellence (NICE), UK advises identifying and managing the underlying causes of AF, treating the arrhythmia and assessing and managing the risk of stroke in these patients.⁹ Clinical prediction scores such as CHA₂DS₂VASc predict the risk of thromboembolic disease including stroke¹⁰ and guide whether the benefits of commencing anticoagulation treatment outweigh risk. The risk of starting a patient on anticoagulation include assessing bleeding risk, for example, using the HAS-BLED score. While components used for calculating CHA₂DS₂VASc and HAS-BLED risk scores are well recorded, risk scores themselves are poorly recorded resulting in a gap in data quality. Anticoagulation therapy aims to reduce the risk of thromboembolic events.

INTRODUCTION

Clinical dashboards integrate large volumes of routine data into a simple accessible format, and are intended to assist clinicians and managers to monitor and improve the quality of care.^{1 2} Dashboards have been used in primary care for a range of functions such as improving data quality and prevention,³ to improve the quality of surveillance⁴ and to promote medication safety.⁵

Evaluation of these dashboards should include a rapid communication on the quality achievement to their target audience. Formal usability studies also assess whether target users of a system interact with it as intended by the designers.⁶ Systematic usability testing and subsequent enhancements increase

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, UK

²Royal College of General Practitioners Research and Surveillance Centre, Royal College of General Practitioners, London, UK

Correspondence to

Professor Simon de Lusignan; simon.delusignan@phc.ox.ac.uk

Figure 1 Screenshot of dashboard section for case ascertainment of incidence/prevalence.

This has been achieved by the use of vitamin K antagonists, primarily warfarin, for many years. The introduction of direct oral anticoagulants (DOACs) such as apixaban and rivaroxaban to clinical practice has changed how AF is managed. DOACs have similar or better mortality and vascular outcomes than warfarin,¹¹ and the added benefit of requiring much less monitoring than warfarin.¹² However, the dosage regime varies between the different DOACs and is complex; errors are common and are associated with hospital admission.¹³ In the UK, the Quality and Outcomes Framework (QoF), a pay-for-performance scheme (P4P), was introduced to provide incentives to incentivise general practitioners to achieve indicator thresholds for managing chronic diseases.¹⁴ This has made a significant improvement to enhancing the quality of AF data being recorded in primary care during the last decade.

We developed an interactive dashboard to provide feedback data quality and the quality of AF management in primary care at the individual general practice level within the Oxford Royal College of General Practitioners (RCGP) Research and Surveillance Centre (RSC) sentinel network. The aim was to provide a tool for general practitioners to monitor data quality on a weekly basis. We

carried out this study to evaluate its usability in primary care.

METHOD

Creating the AF dashboard

We used our generic approach to creating clinical dashboards for a single condition. The use interface requirements and data requirements for the dashboard were developed by practising general practitioners who were members of the study team. This involved identifying data in four sections: (1) Case ascertainment—incidence, prevalence, standardised prevalence and any indicator-related (P4P) prevalence; (2) Indications for therapy and risk factors; (3) Management choices; (4) Quality. We generally avoid more than four to five areas of feedback to avoid overload. The indicators for each section and clinical codes that represent the variables were identified. These clinical codes were used to extract an initial dataset used for developing the dashboard.

We developed the dashboard using Tableau data visualisation software (V.2019.1) which allows data-driven development of dashboards. The initial dashboard was hosted on the public dashboard cloud server and accessed a

Figure 2 Screenshot of dashboard section for calculating stroke and haemorrhage risk.

Figure 3 Screenshot of dashboard section for anticoagulant prescribing.

publicly accessible database server (located within our University IT infrastructure) which hosted only aggregated data to comply with information governance requirements.

Think aloud evaluation

The user interface of a dashboard needs to be evaluated for its usability to ensure a user-friendly and engaging experience. Although questionnaires are the most commonly used method for capturing usability feedback, it has the limitation that feedback is captured after the user interaction has taken place.^{15–18} By contrast, the think aloud provides insight into a system user's cognitive process while carrying out a task.^{19–20} We used the think aloud method to validate the usability of the AF dashboard. During usability testing, study subjects are instructed to verbalise their thoughts while concurrently conducting predefined tasks on the dashboard.²¹

We designed the think aloud session to consist of five tasks. Participants were asked to verbalise their cognitive process while engaging in the tasks and avoid describing the reasons for their actions.²² During the initial four tasks, we asked participant to observe the four main sections of the dashboard. We asked them to observe the given information and interpret with respect to that particular aspect of AF management in their practice. As the fifth task, we asked the participant to observe the complete dashboard and describe the overall state of AF

management in their practice in comparison with the RCGP RSC sentinel network.

Subjects

We invited staff from all general practices participating in the RCGP RSC sentinel network (ie, 320 practices at the time of conducting the study) through the practice newsletter. From those who expressed interest, we invited staff from general practices to cover a range of roles. Participants represented 15 practices located across England. None of the participants were involved in the initial requirement gathering/design of the dashboard. Roles of primary care staff recruited as study participants included general practitioners, nurses and practice managers. We also included clinical researchers and hospital consultants who had expertise in AF as participants. Participants who were not able to attend in person joined using the Gotomeeting remote screen sharing software which also allowed recording screen activity. We aimed to recruit a sample of 20–30 subjects for this study based on guidelines of a previous study.²³

Data capture and analysis

We recorded participants' feedback and screen activity using Gotomeeting screen sharing software (V.10.5). The audio component of the recordings were exported and transcribed by a professional transcription service. The

Figure 4 Screenshot of dashboard section for anticoagulant doses.

Figure 5 Number of visits to the dashboard after the release of the initial and subsequent release of the dashboard enhanced by feedback received during the think aloud study.

transcripts were analysed using NVivo (V.12) qualitative analysis software.

We used grounded theory as our analysis approach. We ensured robustness in our analysis by following a documented logical analysis flow which included two templates. The completed templates were reviewed by a peer researcher to reduce bias of the person conducting the mapping exercise. The following three-step approach was used to analyse the transcripts.

1. Mapping verbalised tasks to sections: Each verbalised task description was extracted and mapped to the corresponding section of the dashboard. We define a 'verbalised task' as the verbal feedback given by the user when performing an interacting with component in the dashboard. The verbalised tasks were mapped to the dashboard sections and similar feedback were grouped (online supplemental table 1).
2. Mapping verbalised tasks to usability problem classes: For each section, we mapped the verbalised tasks to matching usability problem classes. We adapted the usability problem classification method used by Peute *et al* and identified occurrences for each usability problem class.²¹ The usability problem classes include visibility of system status, error messages/help instructions, meaning of labels/graphs, layout/screen organisation and dashboard controls. Furthermore, we classified the identified verbalised tasks based on whether they were positive feedback, negative feedback or sug-

Figure 6 Average number of codes for the different roles of participants across the dashboard sections.

Figure 7 Graphical representation of feedback across the usability problem classes.

gestions for new features (online supplemental table 2).

3. Summarising usability issues across sections/usability problem class: The results table generated by steps 1 and 2 were further condensed to understand sections of the dashboard and type of usability problem which required needed to be addressed.

Enhancing the AF dashboard

The analysis of the dashboard informed which sections required improvements in the user experience. The enhanced dashboard was deployed to general practices in the RCGP RSC sentinel network. We informed general practice staff about these enhancements through user training and updated to dashboard user manuals.

Ethical considerations

Personal data were not collected from the study participants. We obtained informed consent from participant for recording verbal response and screen activities. All participants received oral and written information about the study. We used the HRA decision tool to confirm that no NHS REC ethical approval was required for the study.

RESULTS

AF dashboard

Creating the AF dashboard

We used our generic approach to identify four areas which would give an overview of the quality of AF management: (1) Case ascertainment, (2) Calculating stroke and haemorrhage risk to assess whether the benefits of anticoagulants outweigh their risk,²⁴ (3) Decision to anticoagulate and choice of type, and (4) Prescribing an anticoagulant at the correct dose.²⁵ We developed a dashboard with four sections that corresponded to our generic approach.

Case ascertainment

This section displays the prevalence and incidence of AF within the practice of the participant and this was also compared with practices in the rest of the RCGP RSC sentinel network (figure 1).

Indications for therapy and risk factors

This section displays levels of recording of stroke and haemorrhage risks (CHA₂DS₂VASc scores and HAS-BLED scores) for patients with AF with indication about potential risk score records which can be achieved. NICE guidelines require practitioners to complete CHADVASc scores and HAS-BLED scores for patients with AF (figure 2).

Management choices

Decision to anticoagulate and choice of type is considered in this section of the dashboard. Anticoagulation prescribing levels are given with comparison with overall prescribing in the RCGP RSC sentinel network. All patients with AF with a CHA₂DS₂VASc score more than or equal to 2 should be offered anticoagulation (unless they have an increased bleeding risk) according to NICE guidelines (figure 3).

Quality

This section displays anticoagulation prescribing at the various doses. Different combinations of doses prescribed are displayed for each of the four commonly prescribed four DOACs (figure 4).

Uptake of the AF dashboard in general practice

We hosted the AF dashboard as a part of the MyPractice-Dashboard: a collection of five dashboards that inform participating general practices about performance for different conditions. The access statistics to the MyPracticeDashboard during the period are given in figure 5.

Think aloud evaluation

General results

Professional roles of subjects who participated in the study included general practitioners (n=10), clinical researchers (n=5), practice managers (n=4), nurses (n=4) and pharmacists (n=1) and gender (M=40%, F=60%). Prescription of anticoagulants at suboptimal doses was indicated as the most useful section of the dashboard (57%) although this was also indicated as the most difficult section to interpret of all sections (35%) (figure 6).

Classification of usability feedback

We analysed the content of the feedback according to the three-step method described in Methods section. The smiley faces were considered to be the most effective communicative feature (as indicated by 74% of participants). Thirty per cent of the participants indicated that the dashboard “provided clear feedback” and was “easy to interpret”. Several participants (21%) considered lack of information about the “criteria for the smiley faces” and the “range of smiley faces” as a weakness in the dashboard. A total of 120 verbalised tasks were identified in the 24 transcripts analysed. Individual verbalised tasks categorised as positive, negative and new feature suggestions are given in online supplemental appendix 1. Summarised verbalised tasks mapped to the usability problem classes are given in table 1. We found that the visual representation (figure 7) helpful to interpret the results.

Figure 8 Screenshot of the enhanced version of the risk score section.

Enhancing the AF dashboard

Enhancements to the dashboard were prioritised based on the feedback provided by study participants. The section with risk scores received the most amount of negative feedback. The improved screen layout for this section is given in figure 8. The enhancements included simplifying labelling, limiting numerical information and changing nomenclature according to standards used in other national guidelines (eg, CHA₂DS₂VASc to CHADS₂). We also included links to additional documentation that provided details about how certain values displayed in the section were calculated (eg, calculation of earnings according to the QOF scheme).

The anticoagulation dosing section was enhanced by introducing a simplified taxonomy for anticoagulant dosing which would be more insightful for practice staff (figure 9).

DISCUSSION

Principal findings

This study reported on the think aloud method to assess an AF dashboard. The key finding is that case ascertainment was the section that received most overall positive feedback. Communication of overall performance using the smiley face also received positive feedback. We also found that general practitioners considered the data quality of prescribing section to be most useful to support their work. Furthermore, a key area for improvement was the better annotation of graphs, figures and tables.

While the findings of the evaluation provided an systematic approach for enhancing the dashboard, we

Figure 9 Screenshot of the enhanced version of the anticoagulant doses section based.

Table 1 Number of usability issues (for all participants) across the usability problem classes for the four sections of the dashboard (#1, Case ascertainment; #2, Indications for therapy and risk factors; #3, Management choices; #4, Quality (Prescribing))

Section number	Positive feedback				Negative feedback				New feature suggestions			
	#1	#2	#3	#4	#1	#2	#3	#4	#1	#2	#3	#4
A. Visibility of system status	20	0	2	0	0	2	0	1	2	2	4	1
B. Error messages/help instructions	0	0	0	0	0	0	0	2	0	0	0	5
C. Meaning of labels/graphs	0	11	4	0	0	11	6	8	0	0	0	0
D. Layout/screen organisation	0	0	0	0	8	0	2	6	0	0	3	0
E. Dashboard controls	1	1	0	3	1	4	1	2	0	0	0	3
F. Meaning of tabular data	3	3	0	0	6	0	0	0	1	0	0	0

recognise that there is potential to triangulate the findings by using a heuristic evaluation involving a group of usability experts.²⁶ We were not able to form an expert panel to conduct a heuristic evaluation due to funding and time restrictions.

Implication of findings

The think aloud method of feedback collection provides a systematic approach to prioritise enhancements based on the needs the dashboard users. This helped with increasing adoption rates of dashboard products and in turn has a significant impact towards improving the quality of coded data available for research. As clinical implications, we anticipate that successful adoption of the dashboard will result in improved data quality, resulting in better management of AF in primary care.

Comparison with the literature

Audit-based education methods have been previously used to improve management of association to chronic kidney disease management in primary care.²⁷ Frequent supply of routine data from general practices has allowed continuous data quality monitoring, and this has resulted in improved quality of care and disease surveillance.^{4 28} Similar to other studies that have incorporated user-centred design methods, our enhanced dashboard has successfully demonstrated that understanding user interactions is essential to quality improvement.²⁹

Strengths and limitations

The RCGP RSC provides data quality feedback across a range of other conditions, such as chronic kidney disease and asthma, and strives to maintain consistency in the dashboards used to communicate feedback to practices. Possible prior exposure to some other dashboards may have influenced the results of our think aloud evaluation. We were only able to recruit a small number representing general practitioners. Nevertheless, they constituted 46% of the study group.

The concurrent think aloud approach used for the study has the limitation of being intrusive to the cognitive process over the retrospective think aloud method. This is since the process of providing a simultaneous

commentary while using the dashboard will have an impact on usability.

CONCLUSIONS

We have developed an AF dashboard which has been used for assessing quality of care and reporting feedback to general practices that provide data. We had positive response from the study group that participated in the usability evaluation. The management choices and quality (prescribing) sections of the dashboard were enhanced based on feedback received during the evaluation. Our dashboard appears acceptable to primary care professionals, and such quality improvement interventions should be tested in a trial.

Twitter Harshana Liyanage @harshana

Acknowledgements Practice staff from the participating Royal College of General Practitioners Research and Surveillance Centre (RCGP RSC) practices for contributing to the study. Chris McGee for the development of the dashboard, Hannah McHugh, Zarmina Butt and Noshin Ishrat for supporting the think aloud evaluation.

Contributors SdL conceived the dashboard and formulated this evaluation with substantial contribution from HL. SdL and HL with a substantial contribution from MF and NM drafted the manuscript. JS extracted the data for the dashboard. FF was project manager for this study. All authors contributed to and approve the final version of the paper.

Funding The work was supported by Daiichi Sankyo.

Competing interests RH has received occasional fees from Bayer and Boehringer Ingelheim for speaking or consulting on atrial fibrillation-related stroke risk.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Data can be requested by contacting the corresponding author.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Harshana Liyanage <http://orcid.org/0000-0001-9738-6349>

REFERENCES

- 1 Stadler JG, Donlon K, Siewert JD, *et al.* Improving the efficiency and ease of healthcare analysis through use of data visualization dashboards. *Big Data* 2016;4:129–35.
- 2 Koopman RJ, Kochendorfer KM, Moore JL, *et al.* A diabetes dashboard and physician efficiency and accuracy in accessing data needed for high-quality diabetes care. *Ann Fam Med* 2011;9:398–405.
- 3 McMenamin J, Nicholson R, Leech K. Patient dashboard: the use of a colour-coded computerised clinical reminder in Whanganui regional general practices. *J Prim Health Care* 2011;3:307–10.
- 4 Pathirannehelage S, Kumarapeli P, Byford R, *et al.* Uptake of a dashboard designed to give realtime feedback to a sentinel network about key data required for influenza vaccine effectiveness studies. *Stud Health Technol Inform* 2018;247:161–5.
- 5 Jeffries M, Keers RN, Phipps DL, *et al.* Developing a learning health system: insights from a qualitative process evaluation of a pharmacist-led electronic audit and feedback intervention to improve medication safety in primary care. *PLoS One* 2018;13:e0205419.
- 6 Read A, Tarrell A, Fruhling A. Exploring user preference for the dashboard menu design. *2009 42nd Hawaii International Conference on System Sciences*, IEEE, 2009:1–10.
- 7 Khairat SS, Dukkupati A, Lauria HA, *et al.* The impact of visualization dashboards on quality of care and clinician satisfaction: integrative literature review. *JMIR Hum Factors* 2018;5:e22.
- 8 Schall MC, Cullen L, Pennathur P, *et al.* Usability evaluation and implementation of a health information technology Dashboard of evidence-based quality indicators. *Comput Inform Nurs* 2017;35:281–8.
- 9 Friberg L, Rosenqvist M, Lip GYH. Evaluation of risk stratification schemes for ischaemic stroke and bleeding in 182 678 patients with atrial fibrillation: the Swedish Atrial Fibrillation cohort study. *Eur Heart J* 2012;33:1500–10.
- 10 Chugh SS, Havmoeller R, Narayanan K, *et al.* Worldwide epidemiology of atrial fibrillation: a global burden of disease 2010 study. *Circulation* 2014;129:837–47.
- 11 Ruff CT, Giugliano RP, Braunwald E, *et al.* Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet* 2014;383:955–62.
- 12 Lip GYH. Atrial fibrillation in 2011: stroke prevention in AF. *Nat Rev Cardiol* 2011;9:71–3.
- 13 Bruneau A, Schwab C, Anfosso M, *et al.* Burden of inappropriate prescription of direct oral anticoagulants at hospital admission and discharge in the elderly: a prospective observational multicenter study. *Drugs Aging* 2019;36:1047–55.
- 14 Langdown C, Peckham S. The use of financial incentives to help improve health outcomes: is the quality and outcomes framework fit for purpose? A systematic review. *J Public Health* 2014;36:251–8.
- 15 Schall MC, Cullen L, Pennathur P, *et al.* Usability evaluation and implementation of a health information technology dashboard of evidence-based quality indicators. *Comput Inform Nurs* 2017;35:281–8.
- 16 Dolan JG, Veazie PJ, Russ AJ. Development and initial evaluation of a treatment decision dashboard. *BMC Med Inform Decis Mak* 2013;13:51.
- 17 Daley K, Richardson J, James I, *et al.* Clinical dashboard: use in older adult mental health wards. *Psychiatrist* 2013;37:85–8.
- 18 Martinez W, Threath AL, Rosenbloom ST, *et al.* A patient-facing diabetes dashboard embedded in a patient web portal: design sprint and usability testing. *JMIR Hum Factors* 2018;5:e26.
- 19 Clemmensen T, Hertzum M, Hornbæk K, *et al.* Cultural cognition in the thinking-aloud method for usability evaluation. *Iciss 2008 Proceedings* 2008;1:189.
- 20 Van Someren MW, Barnard YF, Sandberg JA. *The think aloud method: a practical guide to modelling cognitive processes*. London: Academic Press, 1994.
- 21 Peute LWP, de Keizer NF, Jaspers MWM. The value of retrospective and concurrent think aloud in formative usability testing of a physician data query tool. *J Biomed Inform* 2015;55:1–10.
- 22 Ericsson KA, Simon HA. *Protocol analysis: verbal reports as data*. The MIT Press, 1984.
- 23 Boddy CR. Sample size for qualitative research. *Qualitative Market Research: An International Journal* 2016;19:426–32.
- 24 Adeboyeje G, Sylwestrzak G, Barron JJ, *et al.* Major bleeding risk during anticoagulation with warfarin, dabigatran, apixaban, or rivaroxaban in patients with nonvalvular atrial fibrillation. *J Manag Care Spec Pharm* 2017;23:968–78.
- 25 Shum P, Klammer G, Toews D, *et al.* Anticoagulant utilization and direct oral anticoagulant prescribing in patients with nonvalvular atrial fibrillation. *Can J Hosp Pharm* 2019;72:428–34.
- 26 Chen H, Schall MC, Pennathur PR. *Development and evaluation of a health information technology dashboard of quality indicators*. In *Proceedings of the Human Factors and Ergonomics Society annual meeting*. Los Angeles, CA: SAGE Publications, 2015: 59. 461–5.
- 27 Lusignan Sde, de Lusignana S, Gallagher H, *et al.* Audit-based education lowers systolic blood pressure in chronic kidney disease: the quality improvement in CKD (QUICKD) trial results. *Kidney Int* 2013;84:609–20.
- 28 Liyanage H, Williams J, Byford R, *et al.* Near-real time monitoring of vaccine uptake of pregnant women in a primary care sentinel network: ontological case definition across heterogeneous data sources. *Studies in Health Technology and Informatics* 2019;264.
- 29 Colquhoun HL, Sattler D, Chan C, *et al.* Applying user-centered design to develop an audit and feedback intervention for the home care sector. *Home Health Care Manag Pract* 2017;29:148–60.