

Impact of an anticipatory care planning intervention on unscheduled acute hospital care using difference-in-difference analysis

Attakrit Leckcivillize ¹, Paul McNamee,¹ Christopher Cooper,² Robby Steel³

To cite: Leckcivillize A, McNamee P, Cooper C, *et al.* Impact of an anticipatory care planning intervention on unscheduled acute hospital care using difference-in-difference analysis. *BMJ Health Care Inform* 2021;**28**:e100305. doi:10.1136/bmjhci-2020-100305

► Additional online supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjhci-2020-100305>).

Received 17 December 2020
Revised 10 April 2021
Accepted 03 May 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Health Economics Research Unit, University of Aberdeen, Aberdeen, UK

²NHS Lothian Analytical Services, NHS Lothian, Edinburgh, UK

³Division of Psychiatry, Royal Infirmary of Edinburgh, Edinburgh, UK

Correspondence to

Dr Attakrit Leckcivillize; attakrit.leckcivillize@abdn.ac.uk

ABSTRACT

Unscheduled admissions to hospital place great demands on the use of limited healthcare resources in health systems worldwide. A range of approaches exist to manage demand; however, interventions within hospitals have received less attention, and the evidence base on effectiveness is limited. This study aimed to assess the effectiveness of a novel intervention, implemented in National Health Service Lothian, to reduce the number of unscheduled attendances, and to estimate the impact on hospital admissions, length of hospital stay and overall total acute hospital costs.

Methods Before and after observational study of an anticipatory care planning intervention targeted among people identified by a prediction algorithm (Scottish Patients at Risk of Readmission and Admission) as being at high risk of future unscheduled hospital admissions. The statistical significance of the difference in outcomes observed before and after implementation of the intervention between August 2014 and July 2015 was tested using difference-in-difference analysis.

Results The intervention was estimated to reduce the number of unscheduled hospital admissions and emergency department (ED) visits by approximately 0.36 (95% CI –0.905 to 0.191) per patient per year (based on 954 and 450 patients in the intervention and control groups, respectively). There was also non-significant reductions in length of hospital stay for unscheduled admissions and hospital costs for ED visits and inpatient care. The overall predicted effect of the intervention for the average participant was a saving of around £2912 (95% CI –7347.0 to 1523.9) per patient per year.

Conclusion An anticipatory care planning intervention focused among people judged to be at higher risk of future unscheduled hospital admissions can be effective in reducing the number of unscheduled admissions to hospital and ED visits, and may lead to an overall saving in use of hospital resources.

INTRODUCTION

A large focus for healthcare systems worldwide is to improve the efficiency of resource use, especially the amount devoted to unscheduled care in hospital. For example, almost a quarter of National Health Service (NHS) England overall expenditure in

Summary

What is already known?

► Approaches to manage healthcare demand, ranging from provision of self-management support and interventions in primary care, to better integration between healthcare and social care, as well as other specific community-based interventions to reduce hospital re-admission, have been assessed in the literature. Some are beneficial and cost-effective, while others are not, but less is known regarding hospital based interventions.

What does this paper add?

► This paper focused on impacts of an anticipatory care planning intervention based on a prediction algorithm to reduce unscheduled acute hospital care. This approach can be an effective intervention within a hospital to lower unscheduled hospital admissions and emergency department visits, and may potentially reduce overall hospital costs.

2013/2014 was spent on unscheduled care.¹ Unscheduled care has been defined as ‘any healthcare provided with less than 24 hours notice’. Unscheduled hospital admissions form a growing part of hospital care: between 2006/2007 to 2017/2018, the number of unscheduled admissions from accident & emergency (A&E) departments has been steadily rising.²

The UK House of Commons Public Accounts Committee³ estimates that 24% of the total 5.8 million emergency admissions in England during 2016–2017 might have been avoided if more effective community healthcare and case management had existed.⁴ Examples here include interventions co-ordinated in primary care, where one study in NHS Highland showed significant reductions in unplanned hospitalisation for patients with multiple morbidities.⁵ Other initiatives, ranging from self-management support to better



integration between healthcare and social care, have also been tested to reduce hospital readmission (See⁶ for a detailed review). The evidence base relating to the effectiveness of such interventions is however rather mixed, and also challenging to interpret, given patient heterogeneity in terms of case-mix.

Relative to primary and community care, less attention has been devoted to initiatives within the hospital setting. However, there is now growing provision of enhanced information support between healthcare professionals. For example, within Scotland, the key information summary aims to enhance communication between primary and secondary care.⁷ It allows selected parts of the general practitioner (GP) electronic patient record to be shared electronically with other parts of the NHS, and currently covers 2%–3% of the Scottish adult population with the most complex health and/or social care needs.

Predictive modelling tools to identify patients at high risk of being a frequent A&E attender or being readmitted are now available, for instance Patients at Risk of Readmission and Adjusted Clinical Groups-Predictive Model used in the USA and UK,⁸ as well as Hospital Admission Risk Prediction in Canada (see ref. 9 for a rapid review on predictive validity of these tools). In Scotland, an algorithm called Scottish Patients at Risk of Readmission and Admission (SPARRA) is an available risk prediction tool that predicts an individual's risk of unscheduled admission to hospital within the next 12 months. It has been used to proactively manage future hospital demand among population groups likely to make greater use of hospital resources.^{10–11} While modelling studies predict that such tools should lead to reductions in the number of unplanned hospital admissions,¹¹ there is a paucity of evidence from real world implementation studies regarding whether they do in practice reduce the volume and costs associated with unscheduled hospital admissions.¹²

The aim of this paper therefore is to report the effectiveness of a novel intervention, implemented in NHS Lothian, Scotland, that aimed to reduce the risk of future unscheduled hospital admission. The intervention involved application of the SPARRA tool to identify patients at high risk of future unscheduled hospital admission, followed by deployment of appropriate key workers (eg, addictions or psychiatric nurses or consultants) to engage with patients, relatives, GPs and the wider hospital team to develop patient-centred care plans. This paper focuses on four potential effects of the intervention: the number of unscheduled attendances to emergency department (ED) and unscheduled hospital admissions, length of hospital stay, total costs of unscheduled admissions and overall total acute hospital costs. Unlike many interventions that prioritise older adults,^{13–15} this intervention focused on a wider group of participants with long term conditions and younger frequent ED attenders. Therefore, it contributes to the literature on hospital based interventions among a wider patient group.

DATA

The intervention (entitled 'PACT': Patient Anticipatory Care Team) was implemented at the Royal Infirmary Edinburgh (RIE) for 2 years and commenced in August 2014. Frequent hospital attenders were defined as an individual with a SPARRA score of 80 or more, or meeting the conventionally accepted definition of ED frequent attender: 10 or more ED attendances in the previous year or 5 or more in the previous 3 months. Frequent attenders were triaged for interventions and allocated an appropriate PACT key worker, with some patients then receiving anticipatory care plans (ACPs). Staff were able to access this ACP to manage future ED visits.¹⁶ For example, a typical ACP of patients with multiple long-term conditions consisted of background information on physical and mental problems, and family circumstances, for example, carer or care responsibilities. Many included consultant names, as well as flagging specific issues such as need for support from other services, for example, social care or police.

A total of 954 patients were identified as frequent hospital attenders eligible to receive the intervention and were assigned a key worker between August 2014 and July 2015. The algorithm identified a cohort of potentially eligible individuals, which was further refined by removing individuals resident outwith the hospital catchment area or who clinically did not require a care plan (this process also excluded frail elderly patients, who were signposted to dedicated care of the elderly services). Further, for the purposes of analysing full-service outcomes 1 year before and after implementation, only patients alive for the whole outcomes measurement period were included in the statistical analysis (Retaining patients who died during the evaluation period could bias the results in either direction. While greater mortality might drive down the cost due to shorter span of those patients in the sample, end of life care could be much more expensive for some patients.).

In addition, a synthetic control group was created using archived data to identify patients who would have been eligible for the intervention if it had existed on 1 August 2013 (812 patients). After applying the same inclusion criteria, this generated a control group of 450 patients. The flow chart for the triage into intervention and control groups in figures 1 and 2 illustrates the timeline for sample recruitment. The intervention group consists of multiple cohorts of patients identified as high-demand patients at different dates in a monthly staggered pattern. The first cohort of the intervention group was triaged in August 2014. Here, the year 1 (or 'before' period) covers 12 months prior to August 2014, while year 2 (the 'after' period) includes 12 months from August 2014. We combined all cohorts based on triages from August 2014 to July 2015 into 'the intervention group'.




Figure 1 The triage of our study sample.

METHODS

The effectiveness of the intervention was estimated using a ‘difference-in-differences (Dif-in-Difs) design’. The main regression model was specified as follows:

$$y_{it} = \alpha + \beta Treat_{it} + \delta After_{it} + \gamma Treat_{it} * After_{it} + \theta X_{it} + \epsilon_{it}$$

where y_{it} is a dependent variable for the outcomes of interest, which are numbers of ED visits, unscheduled hospital admissions, length of stay or total hospital cost as described in [table 1](#). $Treat_{it}$ is a dummy variable equal to one if patient i was allocated to the intervention, $After_{it}$ is a dummy variable equal to one if the time period t ($=1, 2$) of the data is within 1 year after patient i was allocated to the intervention and 1 year after the hypothetical intervention date (1 August 2013) for the intervention and control groups respectively, and γ , a coefficient of $Treat_{it} * After_{it}$, is the estimated impact of the intervention on the outcomes (treatment effect). X_{it} is a vector of control variables including age, gender, SPARRA score when identified by the algorithm as a frequent attender, and quintile of the patient’s neighbourhood based on the 2016 Scottish Index of Multiple Deprivation (SIMD), where the first quintile is the most deprived area whereas the fifth quintile is the least deprived area. ϵ_{it} is an error term. There were a small number of missing observations in the SPARRA score and the SIMD ($n=154$). Thus, the

data were treated as missing at random and therefore those observations were excluded from the main analysis. Negative binomial regression was used to model the number of hospital admissions and length of stay, while generalised linear model regression with gamma distribution and logarithmic link function was used for total hospital costs.

RESULTS

[table 1](#) shows per patient mean, SD, minimum and maximum value of the main outcomes and all explanatory variables.

The average number of unscheduled admissions and visits to the ED was 3.86 per person per year (pp/py), of which the ED visit is 0.95 and the unscheduled admission is 2.91. All values of number of admissions/visits presented in [tables 1 and 2](#) as well as in the regression analysis of total cost in online supplemental table 1 and 2 are calculated assuming one admission/visit per episode of care, for example, when a patient visited the ED and was then transferred to another ward, we count that as one admission/visit only (using a different approach, ie, counting them as separate admissions or visits, makes little difference to the regression analysis results.). The




Figure 2 Timeline of the administrative data used for the control and intervention groups.

**Table 1** Summary statistics

Variable	Obs	Mean (N, proportion)	SD	Min	Max
No of unscheduled hospital admissions and ED visits (visits/person/year)	2808	3.86	3.97	0	33
No of unscheduled hospital admissions (non-ED) (admissions/person/year)	2808	2.91	3.25	0	31
No of ED visits (visits/person/year)	2808	0.95	1.43	0	14
Length of stay for unscheduled admissions (days)	2808	9.43	15.29	0	290
Total cost from unscheduled admissions and ED visits (£/person/year)	2808	10829	14550	0	284252
Total cost for ED visits+scheduled and unscheduled admissions (£/person/year)	2808	22249	32287	0	294837
Female	2808	(1,374, 48.9%)	–	0	1
Age group 16–24	2808	(266, 9.5%)	–	0	1
Age group 25–34	2808	(418, 14.9%)	–	0	1
Age group 35–44	2808	(464, 16.5%)	–	0	1
Age group 45–54	2808	(608, 21.7%)	–	0	1
Age group 55–64	2808	(512, 18.2%)	–	0	1
Age group 65–74	2808	(478, 17.0%)	–	0	1
Age group 75 up	2808	(62, 2.2%)	–	0	1
SIMD quintile 1	2712	(810, 29.9%)	–	0	1
SIMD quintile 2	2712	(872, 32.2%)	–	0	1
SIMD quintile 3	2712	(418, 15.4%)	–	0	1
SIMD quintile 4	2712	(306, 11.3%)	–	0	1
SIMD quintile 5	2712	(306, 11.3%)	–	0	1
SPARRA score at entry	2676	70.27	19.78	6	95

ED, emergency department; SIMD, Scottish Index of Multiple Deprivation; SPARRA, Scottish Patients at Risk of Readmission and Admission.

average length of stay for unscheduled hospital admissions is 9.4 days pp/py. The average total cost for unscheduled hospital admissions is £10 830 pp/py, and the average total cost for all ED visits and all admissions is £22 250. Regarding explanatory variables, there are approximately the same proportion of males and females in the sample. Approximately 40% of the sample is aged between 16–44 years of age, 40% aged 45–64 years of age, and those aged 65 years of age or older represent 20% of the sample. Approximately 60% of the sample live in less privileged neighbourhoods (quintiles 1 and 2), as measured by SIMD. The average baseline SPARRA score is 70.

Table 2 compares descriptive statistics for the explanatory variables: age, gender, SIMD and SPARRA between the control and intervention groups. The groups appear comparable on most indicators, with no large differences evident. Summary statistics of the four main outcome variables are also shown. The average number of admissions and ED visits show a clear downward trend between the before and after period for both the intervention group and the control group (average decline of 1 visit pp/py in both groups). The average length of stay associated with unscheduled admissions also shows a decline in both groups, with a smaller decline in the intervention

group (3 days) vs the control group (4 days) (However, these are not adjusted for the compositional differences (such as age or risk score) between the intervention and control groups.). A similar pattern emerges with the costs associated with these stays: a decline in both groups, but a larger decline in the control group. Finally, the total costs from all ED visits and all hospital admissions showed a similar level decline for both intervention and control groups between the before and after period.

REGRESSION ANALYSIS

Table 3 shows the intervention group has a significantly lower number of unscheduled admissions and ED visits. Also, there were significantly fewer admissions and ED visits in the period after intervention. The effect of the intervention is shown by the interaction term (intervention*after). This estimates a reduction in the number of unscheduled admissions and ED visits, and is statistically significant at the 10% level. This result is robust to a change in the method used to correct the standard errors, that is, the p value of the treatment effect is equal to 0.070 and 0.085 for homoscedastic errors and bootstrapping errors with stratification between control and

Table 2 Summary statistics by group and period

Variable	Intervention group					Control group				
	Obs	Before		After		Obs	Before		After	
		Mean	SD	Mean	SD		Mean	SD	Mean	SD
No of unscheduled hospital admissions & ED visits	954	4.09	3.62	2.89	3.70	450	5.19	3.99	4.06	4.67
No of unscheduled hospital admissions (non-ED)	954	3.10	3.06	2.16	2.91	450	3.91	3.34	3.12	3.84
No of ED visits	954	1.01	1.31	0.73	1.40	450	1.29	1.59	0.95	1.47
Length of stay for unscheduled admissions	954	9.87	14.41	6.83	12.43	450	13.68	20.75	9.77	15.15
Total cost from unscheduled admissions and ED visits only	954	11 110	13 057	7 455	10 876	450	16 447	20 127	11 771	15 769
Total cost for ED visits+scheduled and unscheduled admissions	954	22 822	30 329	16 763	28 632	450	30 702	36 865	24 218	36 409
Personal characteristics*										
Female	954	0.51	0.50			450	0.44	0.50		
Age group 16–24	954	0.09	0.29			450	0.10	0.31		
Age group 25–34	954	0.16	0.37			450	0.12	0.33		
Age group 35–44	954	0.17	0.37			450	0.16	0.37		
Age group 45–54	954	0.22	0.42			450	0.21	0.41		
Age group 55–64	954	0.16	0.37			450	0.23	0.42		
Age group 65–74	954	0.17	0.38			450	0.17	0.37		
Age group 75 up	954	0.03	0.17			450	0.00	0.07		
SIMD Quintile 1	913	0.29	0.45			443	0.32	0.47		
SIMD Quintile 2	913	0.31	0.46			443	0.35	0.48		
SIMD Quintile 3	913	0.16	0.37			443	0.13	0.34		
SIMD Quintile 4	913	0.12	0.32			443	0.10	0.31		
SIMD Quintile 5	913	0.12	0.32			443	0.10	0.30		
SPARRA score at entry	914	68.21	20.14			424	74.72	18.22		

*We have access to the personal characteristics of the patients only once at the time they entered the intervention group (or the hypothetical intervention date for the control group), hence we do not present the numbers for the ‘after’ period. ED, emergency department; SIMD, Scottish Index of Multiple Deprivation; SPARRA, Scottish Patients at Risk of Readmission and Admission.

intervention group, respectively. Most of the remaining control variables in [table 3](#) have insignificant coefficients. The exceptions relate to the youngest age group, SIMD 1 and the SPARRA score.

To interpret the estimated coefficients shown in [table 3](#), marginal effects and CI are computed, based on the delta method. [table 4](#) shows the predicted number of unscheduled admissions and ED visits per year in the intervention group is around 3.30 compared with 3.86 in the control group, while the number in the period after intervention falls by around 0.91. In summary the intervention is associated with a reduction of approximately 0.357 (95% CI –0.905 to 0.191) admissions and visits per year (This is approximately equal to (2.8424–3.8477) – (3.5538–4.2023)). The treatment effect and 95% CI are computed according to a suggestion in reference [17](#). Further, the predicted effects of the intervention may be larger for those with higher SPARRA scores or younger patients (see online supplemental table A1).

[table 5](#) shows the impact on hospital length of stay from unscheduled admissions. There was no statistically significant difference in length of stay between intervention and control groups, or in the time period before and after the intervention. The interaction term (intervention*after) is not statistically significant, suggesting that the intervention did not influence length of stay. Female patients were associated with a longer length of stay. Those from the most deprived areas were associated with a shorter length of stay (Although this result sounds counter-intuitive, this coefficient becomes significant only when SPARRA score is included. In other words, for those with the same risk score, patients from the most deprived area stay in the hospital for shorter periods of time than those from the least deprived area.). Higher SPARRA score was significantly associated with a longer length of stay. Overall, while the ‘average’ patient is predicted to have a shorter length of stay by approximately 1 day, the effect is not statistically different from zero.



Table 3 Negative binomial regression with bootstrapped SEs (1000 replications) without stratification for unscheduled admissions and ED visits*

Variable	Coef.	SE	Z-stat	P value	(95% CI)
Intervention	-0.0882	0.0433	-2.04	0.042	-0.1731 to -0.0033
After	-0.1676	0.0634	-2.64	0.008	-0.2919 to -0.0433
Intervention*after	-0.1352	0.0790	-1.71	0.087	-0.2901 to 0.0197
Female	-0.0047	0.0387	-0.12	0.902	-0.0806 to 0.0711
Age 16–24	0.2696	0.1194	2.26	0.024	0.0355 to 0.5037
Age 25–34	0.1821	0.1075	1.69	0.09	-0.0286 to 0.3928
Age 35–44	0.1436	0.1113	1.29	0.197	-0.0745 to 0.3616
Age 45–54	-0.1244	0.1023	-1.22	0.224	-0.3248 to 0.0761
Age 55–64	-0.1626	0.1066	-1.52	0.127	-0.3716 to 0.0464
Age 65–74	-0.0749	0.1082	-0.69	0.489	-0.2869 to 0.1371
Age 75 up					Reference group
SIMD quintile 1	-0.1573	0.0726	-2.17	0.03	-0.2996 to -0.0151
SIMD quintile 2	-0.0017	0.0723	-0.02	0.981	-0.1434 to 0.1400
SIMD quintile 3	-0.0221	0.0812	-0.27	0.786	-0.1813 to 0.1372
SIMD quintile 4	0.0916	0.0864	1.06	0.289	-0.0778 to 0.2610
SIMD quintile 5					Reference group
SPARRA score at entry	0.0254	0.0015	17.35	0	0.0226 to 0.0283
Intercept	-0.3125	0.1685	-1.85	0.064	-0.6427 to 0.0177

Number of obs=2654.

Log likelihood=-6247.4205 Pseudo R2=0.0455.

*The results in tables 3 and 4 allow for multiple admissions per day in order to reflect resources and staff time used with each patient. However, the regression results based on number of admissions capped at one admission per day are both qualitatively and quantitatively similar to tables 3 and 4.

ED, emergency department; SIMD, Scottish Index of Multiple Deprivation; SPARRA, Scottish Patients at Risk of Readmission and Admission.

Online supplemental table 1 estimates the association between the intervention and unscheduled hospital admission costs. The coefficient of the interaction variable— (intervention* after)—is not significantly different from zero, indicating no statistically significant association with unscheduled admission costs. Similar to previous tables, gender, area deprivation and SPARRA score were all significantly associated with costs. Using the estimated coefficients, our model predicts a £717 (95% CI -2752.1

to 1318.5) reduction in unscheduled hospital admission costs for the average patient.

Online supplemental table 2 considers the effect on total hospital costs for ED visits and all lengths of stay (both scheduled and unscheduled). The coefficient of interest (intervention*after) is not significantly different from zero. Similar to previous estimates, gender, area deprivation and SPARRA score were all significantly associated with costs (male respondents, younger age groups,

Table 4 Predicted number of unscheduled admissions estimated at means of all other control variables based on bootstrapped SE without stratification

	Predicted value	Delta-meth. SE	(95% CI)
Control group	3.8645	0.1341	3.6017 to 4.1273
Intervention group	3.3071	0.0831	3.1443 to 3.4699
Before	3.9566	0.0850	3.7901 to 4.1231
After	3.0506	0.1036	2.8475 to 3.2537
Control and before	4.2023	0.1535	3.9014 to 4.5033
Control and after	3.5538	0.1974	3.1670 to 3.9407
Intervention and before	3.8477	0.0981	3.6554 to 4.0401
Intervention and after	2.8424	0.1187	2.6098 to 3.0750

Table 5 Negative binomial regression with bootstrapped SEs (1000 replications) without stratification for length of stay due to unscheduled admissions

Variable	Coef.	SE	Z-stat	P value	(95% CI)
Intervention	-0.1035	0.0746	-1.39	0.165	-0.2498 to 0.0428
After	-0.1506	0.0983	-1.53	0.125	-0.3432 to 0.0420
Intervention*after	-0.1332	0.1240	-1.07	0.283	-0.3761 to 0.1098
Female	0.2300	0.0625	3.68	0	0.1075 to 0.3525
Age 16–24	-0.0818	0.1856	-0.44	0.659	-0.4456 to 0.2820
Age 25–34	-0.0675	0.1784	-0.38	0.705	-0.4171 to 0.2822
Age 35–44	0.0033	0.1758	0.02	0.985	-0.3414 to 0.3479
Age 45–54	-0.1204	0.1719	-0.7	0.484	-0.4573 to 0.2165
Age 55–64	-0.0176	0.1729	-0.1	0.919	-0.3564 to 0.3213
Age 65–74	0.1484	0.1714	0.87	0.387	-0.1875 to 0.4844
Age 75 up					Reference group
SIMD Quintile 1	-0.3193	0.1156	-2.76	0.006	-0.5460 to -0.0927
SIMD Quintile 2	-0.0892	0.1150	-0.78	0.438	-0.3146 to 0.1362
SIMD Quintile 3	-0.2530	0.1219	-2.08	0.038	-0.4920 to -0.0141
SIMD Quintile 4	0.1089	0.1312	0.83	0.406	-0.1482 to 0.3661
SIMD Quintile 5					Reference group
SPARRA score at entry	0.0345	0.0023	15.12	0	0.0301 to 0.0390
Intercept	-0.1317	0.2710	-0.49	0.627	-0.6629 to 0.3995

Number of obs=2654.

Log likelihood=-8168.7977 Pseudo R2=0.0324.

SIMD, Scottish Index of Multiple Deprivation; SPARRA, Scottish Patients at Risk of Readmission and Admission.

more deprived area and lower SPARRA score were all associated with lower hospital costs).

We also ran bootstrap regressions of the model in online supplemental table 2 for 10000 replications and plotted the histogram of the estimated coefficients of the intervention effects in online supplemental figure A1. The average value of these estimates is around -0.18, and in more than 90% of the replications the intervention effects are negative, showing a reduction in total costs from all scheduled and unscheduled admissions as well as ED visits.

Online supplemental table 3 presents the marginal effects for different groups. For example, the predicted value of the total cost per year in the intervention group is around £17230 compared with £19985 for the control group. Furthermore, for both groups, the total cost observed after the intervention period is around £3230 lower than observed in the year prior to intervention. Our predicted effect of the intervention for the 'average' patient is a saving of around £2912 (95% CI -7347.0 to 1523.9) per patient per year. Online supplemental table A2 indicates that the predicted savings are likely to be larger among those patients with higher SPARRA score, for example, £4484 (95% CI -11 309.7 to 2342.8) and £5133 (95% CI -12 948.9 to 2683.6) pp/py for those with SPARRA score at 75th and 90th percentile respectively. In addition, the absolute size of the reductions is increasing with age except for the oldest patients, while

female patients are predicted to have a higher decrease in costs than males.

DISCUSSION

The results reported in this paper suggest that the ACP intervention developed in this study had a statistically significant association with the number of unscheduled hospital admissions and ED visits. However, while the intervention was associated with a reduced length of hospital stay for unscheduled admissions, and was associated with lower hospital costs for ED visits and inpatient care, these reductions were not statistically significant. Such insignificant effects are consistent with literature on case management in older people and care pathway guidelines for specific diseases.^{13 18}

Our finding of a statistically significant reduced volume of unscheduled hospital admissions and ED visits is consistent with systematic reviews of similar interventions that aim to reduce hospital readmissions (19,20). A meta-analysis of 42 RCTs (Randomised Controlled Trials)¹⁹ found a significant reduction in the risk of 30 days medical and surgical readmissions (pooled relative risk=0.82 (95% CI 0.73 to 0.91)). The ingredients within successful interventions targeted patient characteristics (eg, multimorbidity and socioeconomic variables factors) as well as caregiver availability. A further systematic review of 26 RCTs²⁰ found that interventions initiated during

the hospital stay and continuing after discharge for a minimum of 1 month were effective in reducing readmissions over the subsequent 1 year period (pooled relative risk=0.58 (95% CI 0.46 to 0.75)).

One explanation for the insignificant findings on three of the four outcomes is that there was a clear downward in the second year of follow-up in both groups. This is most likely due to regression to the mean, that is, the trajectory of hospital care use reverting to a trend that reflects more typical longer term use patterns. The use of the SPARRA algorithm and prior admission data will by design detect high users of hospital care. One implication is that reductions in admissions and length of stay are more likely to be difficult to detect, and may lead to zero effects, or weakly significant treatment effects.

An explanation for the insignificant effect of the intervention on costs is that the intervention is focused on prevention of admission rather management of length of stay while in hospital. A further explanation is that length of stay may be influenced by bed capacity, and this may have fluctuated year on year, thereby diluting any treatment effect.

A potential limitation of this study is that the estimates relied on historical control data. Other than standard Dif-in-Difs assumptions (Ideally, a better control group could be contemporaneous data from other health boards that did not implement the PACT intervention. However, we do not have access to data outside the RIE.), one extra assumption is needed when using patients' historical data as a control group. Specifically, the change in outcomes of patients in the control group 1 year before and after 1 August 2013 (our hypothetical intervention date) is assumed to mimic what would have happened to patients in the intervention group if they had not been treated with the intervention (similar to the 'parallel trend' assumption).

However, we encounter two complications. First, due to data availability, we are unable to show that the trends between the intervention and control groups were similar prior to the actual and hypothetical dates respectively. Second, the dummy $After_{it}$ is not able to control for macro trends affecting both intervention and control groups in the year after intervention because these periods are not contemporaneous. Consequently, the analysis assumes no significant change in resource constraints between different time periods. Overall, these limitations and the unavailability of data on other patients' characteristics, may limit the generalisability of our findings to other settings. Further research in other settings is required to establish whether the results observed here can be generalised to other areas.

CONCLUSIONS

An anticipatory care planning intervention focused on people at high risk of future unscheduled hospital admissions can be effective in lowering the number of

unscheduled admissions to hospital and ED visits, and may help to save overall hospital costs. Furthermore, potential benefits of the intervention in terms of health for the patient were not measured. These benefits may include better health outcomes and positive patient experience.²¹ Although feedback from patients, relatives and staff was overwhelmingly positive,¹⁶ evaluation of the intervention in terms of these aspects was beyond the scope of this study.

Acknowledgements We thank Information Statistics Division and the data analytical team from NHS Lothian for data linkage.

Contributors AL performed data analysis, undertook data interpretation and wrote drafts of the manuscript. PM undertook data interpretation, wrote drafts of the manuscript and edited the manuscript prior to publication. CC helped with data preparation, data linkage and organised information governance permissions. RS led the original study on which the paper is based and contributed to the writing of the manuscript. All authors read and approved the final manuscript.

Funding This work was supported by NHS Lothian through an award from the Invest to Save Fund. The Health Economics Research Unit is funded by the Chief Scientist Office of the Scottish Government Health and Social Care Directorates.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethics approval and consent to participate was obtained from the NHS Lothian Caldicott Guardian (Application Number 1751).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. Data access has to be applied with the NHS Lothian Caldicott Guardian.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Attakrit Leckcivizile <http://orcid.org/0000-0002-6172-3902>

REFERENCES

- 1 NHS England. 2013/14 CCG programme budgeting benchmarking tool. [Internet]. London, 2015. Available: <https://www.england.nhs.uk/wp-content/uploads/2015/06/13-14-ccg-prog-bug-benchmarking-tool.xlsm> [Accessed 30 Oct 2019].
- 2 Steventon A, Deeny S, Friebe R. Emergency hospital admissions in England: which may be avoidable and how?. Briefing Paper. [Internet]. London; The Health Foundation, 2018. Available: <https://www.health.org.uk/publications/emergency-hospital-admissions-in-england-which-may-be-avoidable-and-how> [Accessed 20 Aug 2019].
- 3 House of Commons Committee of Public Accounts. Reducing emergency admissions. [Internet]. London; June, 2018. Available: <https://publications.parliament.uk/pa/cm201719/cmselect/cmpubacc/795/79502.htm> [Accessed 20 Aug 2019].
- 4 Torjesen I. Almost 1.5m emergency hospital admissions could have been avoided last year. *BMJ* 2018;361:k2542.

- 5 Baker A, Leak P, Ritchie LD, *et al*. Anticipatory care planning and integration: a primary care pilot study aimed at reducing unplanned hospitalisation. *Br J Gen Pract* 2012;62:e113–20.
- 6 Purdy S. Avoiding hospital admissions. what does the research evidence say?. London; King's Fund, 2010. Available: https://www.kingsfund.org.uk/sites/default/files/Avoiding-Hospital-Admissions-Sarah-Purdy-December2010_0.pdf [Accessed 18 Aug 2019].
- 7 Hall CC, Lunan C, Finucane A, *et al*. Improving access to the KIS in secondary care. *BMJ Open Qual* 2017;6:e000114.
- 8 Woodhams V, de Lusignan S, Mughal S, *et al*. Triumph of hope over experience: learning from interventions to reduce avoidable hospital admissions identified through an academic health and social care network. *BMC Health Serv Res* 2012;12:153.
- 9 Paton F, Wilson P, Wright K. Predictive validity of tools used to assess the risk of unplanned admissions: a rapid review of the evidence. Evidence Briefings, 2014. Available: <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.570.6517&rep=rep1&type=pdf> [Accessed 25 Mar 2021].
- 10 Mahmoud A. Scottish patients at risk of readmission and admission (Sparra). *Int J Integr Care* 2016;16:216.
- 11 Wallace E, Smith SM, Fahey T, *et al*. Reducing emergency admissions through community based interventions. *BMJ* 2016;352:h6817.
- 12 Marple J. Reducing unplanned hospital admissions using an electronic system for sharing anticipatory care plans between primary and secondary care. *Int J Integr Care* 2016;16:229.
- 13 Huntley AL, Thomas R, Mann M, *et al*. Is case management effective in reducing the risk of unplanned hospital admissions for older people? A systematic review and meta-analysis. *Fam Pract* 2013;30:266–75.
- 14 Wallace E, Stuart E, Vaughan N, *et al*. Risk prediction models to predict emergency hospital admission in community-dwelling adults: a systematic review. *Med Care* 2014;52:751–65.
- 15 Walsh B. Unplanned admissions and readmissions in older people: a review of recent evidence on identifying and managing high-risk individuals. *Rev Clin Gerontol* 2014;24:228–37.
- 16 The Scottish Government. Personalising realistic medicine: chief medical officer for Scotland's annual report 2017-2018. [Internet]. Edinburgh, 2019. Available: <https://www.gov.scot/publications/personalising-realistic-medicine-chief-medical-officer-scotland-annual-report-2017-2018> [Accessed 18 Aug 2019].
- 17 Puhani PA. The treatment effect, the cross difference, and the interaction term in nonlinear “difference-in-differences” models. *Econ Lett* 2012;115:85–7.
- 18 Purdy S, Paranjothy S, Huntley A. Interventions to reduce unplanned hospital admission: a series of systematic reviews. Final Report. [Internet]. Edinburgh, 2012. Available: <http://www.bristol.ac.uk/media-library/sites/primaryhealthcare/migrated/documents/unplannedadmissions.pdf> [Accessed 30 Oct 2019].
- 19 Leppin AL, Gionfriddo MR, Kessler M, *et al*. Preventing 30-day Hospital readmissions: a systematic review and meta-analysis of randomized trials. *JAMA Intern Med* 2014;174:1095–107.
- 20 Verhaegh KJ, MacNeil-Vroomen JL, Eslami S, *et al*. Transitional care interventions prevent Hospital readmissions for adults with chronic illnesses. *Health Aff* 2014;33:1531–9.
- 21 Ross S, Goodwin N, Curry N. Case management: what is it and how it can best be implemented. London; King's Fund, 2011. Available: https://www.kingsfund.org.uk/sites/default/files/Case-Management-Paper-The-Kings-Fund-Paper-November-2011_0.pdf [Accessed 18 Aug 2019].