


User testing of a diagnostic decision support system with machine-assisted chart review to facilitate clinical genomic diagnosis

Alanna Kulchak Rahm ¹, Nephi A Walton,² Lynn K Feldman,³ Conner Jenkins,⁴ Troy Jenkins,⁴ Thomas N Person,¹ Joeseeph Peterson,⁴ Jonathon C Reynolds,¹ Peter N Robinson,^{5,6} Makenzie A Woltz,¹ Marc S Williams,¹ Michael M Segal³

To cite: Kulchak Rahm A, Walton NA, Feldman LK, *et al.* User testing of a diagnostic decision support system with machine-assisted chart review to facilitate clinical genomic diagnosis. *BMJ Health Care Inform* 2021;**28**:e100331. doi:10.1136/bmjhci-2021-100331

► Additional online supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjhci-2021-100331>).

Received 25 January 2021
Revised 01 April 2021
Accepted 20 April 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Alanna Kulchak Rahm;
akrahm@geisinger.edu

ABSTRACT

Objectives There is a need in clinical genomics for systems that assist in clinical diagnosis, analysis of genomic information and periodic reanalysis of results, and can use information from the electronic health record to do so. Such systems should be built using the concepts of human-centred design, fit within clinical workflows and provide solutions to priority problems.

Methods We adapted a commercially available diagnostic decision support system (DDSS) to use extracted findings from a patient record and combine them with genomic variant information in the DDSS interface. Three representative patient cases were created in a simulated clinical environment for user testing. A semistructured interview guide was created to illuminate factors relevant to human factors in CDS design and organisational implementation.

Results Six individuals completed the user testing process. Tester responses were positive and noted good fit with real-world clinical genetics workflow. Technical issues related to interface, interaction and design were minor and fixable. Testers suggested solving issues related to terminology and usability through training and infobuttons. Time savings was estimated at 30%–50% and additional uses such as in-house clinical variant analysis were suggested for increase fit with workflow and to further address priority problems.

Conclusion This study provides preliminary evidence for usability, workflow fit, acceptability and implementation potential of a modified DDSS that includes machine-assisted chart review. Continued development and testing using principles from human-centred design and implementation science are necessary to improve technical functionality and acceptability for multiple stakeholders and organisational implementation potential to improve the genomic diagnosis process.

INTRODUCTION

Clinical decision support (CDS) integrated into electronic health records (EHRs) has long been considered a promising way to improve patient outcomes and decrease inefficiencies.^{1–4} It is also recognised that CDS

Summary

What is already known?

- There is a need in clinical genomics for tools that assist in analysis of genomic information and can do so using information from the electronic health record.
- Such tools should be easy to use, fit within clinical workflows, and provide solutions to priority problems as defined by clinician end-users.
- Natural language processing (NLP) is a useful tool to read patient records and extract findings.

What does this paper add?

- We demonstrated the use of Human-centred design and implementation science principles in a simulated environment for assessment of a new version of a decision support tool prior to large-scale implementation.
- This study provides preliminary evidence that a clinical decision support tool with machine-assisted chart review is acceptable to clinical end-users, fits within the clinical workflow, and addresses perceived needs within the differential diagnosis process across all Mendelian genetic disorders.
- Terminology codes for diagnostic decision support systems should have levels of granularity tuned to the sensitivity and specificity appropriate to its various functions, for example, NLP versus chart documentation.

must be designed with the user in mind, fitting the concepts of human-centred design with computer interfaces at the individual clinician level.^{1–5} Design alone, however, is insufficient to facilitate implementation. For CDS to impact care and patient outcomes, it must fit within clinician workflow and provide a solution to a priority problem for the clinician and the healthcare system.^{4–8}

Diagnostic decision support systems (DDSSs) are a key type of CDS needed in genomics to supplement a shortage of

trained clinicians and address the inherent complexity of genomic diagnosis.^{9 10} This complexity arises from the heterogeneous nature of genetic diseases, the variable expression in patients and the degree of overlap in findings (ie, signs, symptoms and test results) among genetic conditions, sometimes differentiated only by onset age of individual findings.¹¹ Position statements and a systematic review note two new functions needed for DDSSs in genomics: (1) a cost-effective, regular approach to re-evaluation of patient cases in light of new findings or genetic knowledge, when testing does not immediately yield a diagnosis; and (2) developing machine-assisted chart review.^{12 13} Most genomic patient records are extensive with input from by multiple clinicians, such that manual review is prohibitively time-consuming; resulting in added costs from repeated or unnecessary tests and increased risk of missed information that could have facilitated timely diagnosis. Because most of the relevant information is in unstructured clinical notes, approaches such as natural language processing (NLP) are needed to automate and assist this manual process.

To address both re-evaluation and automation, we adapted a commercially available DDSS already capable of incorporating genomic sequencing data to perform automated chart review and present the information to a clinician in the form of findings obtained through structured data mining and NLP of an EHR. We then created clinical case vignettes to simulate the real-world clinical diagnostic workflow for user testing. The goal was to provide preliminary evidence of usability, perceived fit with clinical need and workflow, and potential for implementation into the real-world clinical environment.

METHODS

Setting

Development of the clinical case vignettes, simulated EHR environment, and user testing were conducted at Geisinger, a healthcare system in rural Pennsylvania.

Adapting a DDSS for machine-assisted chart review of clinical findings

We adapted SimulConsult's Genome-Phenome Analyzer, as it is the one DDSS that allows for detailed analysis of clinical information, including pertinent negatives, findings onset information and frequency and treatability of diseases. It has also been shown to be accurate and helpful in clinical diagnosis, including interpreting genomic results.^{14–16} Described in detail elsewhere,^{11 14 15} SimulConsult correlates annotated variant call files (VCFs) with patient-specific clinical and family history information; and the underlying algorithms include age-dependent Bayesian pattern-matching and computational metrics of usefulness and pertinence. SimulConsult also generates a Patient Summary for saving interim patient findings and a customisable genomic return of results (RoR) report shown in previous research to be effective for facilitating standardised communication for patients and referring clinicians.^{17–20} When clinicians enter findings, the DDSS returns a ranked list of candidate diseases and suggestions of other findings to check, ranked by usefulness in narrowing the differential diagnosis in a way that accounts for cost and treatability; thus facilitating the iterative approach of information gathering in diagnosis.^{21 22} For each finding, the presence (with onset age) or absence can be specified (figure 1).

We used the Logica platform to create a simulated EHR and the cTAKES tool with the Unified Medical Language




Figure 1 SimulConsult main interface showing ranked list of candidate diseases and guidance for entering finding presence (or absence) with onset age.

Table 1 Adaptations made to existing DDSS to create GPACSS

Adaptation	Component	Approach
Overall design	SMART-on-FHIR enabled EHR	► Logica platform (https://www.logicahealth.org/ ; formerly Health Services Platform Consortium)
	Archive	► Custom archive stores key files ► RESTful interface.
Coordination and communication	User interface	► SMART-on-FHIR application (GPACSS FHIR app client, figure 2). ► Interface allows user access to DDSS directly from patient record. ► Choice to launch with no findings or with findings previously saved.
	Coordination	► GPACSS ‘Coordinator’ application programming interface (API) saves the NLP output ► Matching of UMLS codes in NLP output to DDSS findings ► Send the matched flagged findings to the DDSS at launch (figure 2)
Natural language processing	Extraction of findings	► NLP: open source Apache cTAKES V.4.0. ²³ ► cTAKES default modules to handle sentence boundary detection, tokenisation, normalisation, tagging parts of speech, recognising named entities and negation. ► cTAKES pretrained module to recognise UMLS concepts in text.
	Mapping in DDSS	► DDSS findings mapped within the DDSS to one or more UMLS and Human Phenotype Ontology codes. ► Mapping strategy minimises false negatives in term capture while tolerating false positives (identifying information unrelated or irrelevant to the diagnostic process).
	Display in DDSS	► Findings identified by NLP display a flag icon. ► Clicking the flag enables viewing of metadata.

DDSS, diagnostic decision support system; EHR, electronic health record; GPACSS, Genotype-Phenotype Archiving and Communication System with SimulConsult; UMLS, Unified Medical Language System.

System (UMLS) module²³ for NLP of patient notes. Steps in adaption included (1) mapping DDSS findings to Human Phenotype Ontology (HPO) and UMLS codes, including creation of hundreds of new HPO terms resulting in creation of new UMLS concepts, (2) using results from NLP analysis of EHR notes to flag ‘Mentions’ of the findings used by the DDSS and (3) augmenting the DDSS’s interface to present the flagged findings with contextual information needed to clinically assess the information ([table 1](#)).

The architecture of the resulting prototype, called the Genotype-Phenotype Archiving and Communication System with SimulConsult (GPACSS), is shown in [figure 2](#).

Clinician review of the flagged findings created from the automated findings search using NLP is facilitated through flag icons ([figure 3](#)). Through this ‘machine-assisted’ chart review, the clinician reviews flagged findings and decides whether and how to specify presence (with a particular onset) or absence (or omit) as shown in [figure 1](#). The mapping of DDSS findings to multiple




Figure 2 Architecture of the Genotype-Phenotype Archiving and Communication System with SimulConsult (GPACSS). The key components are the coordination/archiving system, the DDSS and the NLP. DDSS, diagnostic decision support system; EHR, electronic health record; NLP, natural language processing.




Figure 3 Flagged findings with EHR text display for DDSS. A finding having a flag icon indicates that information was found in the EHR. Clicking the flag shows the various mentions of the flagged finding. DDSS, diagnostic decision support system; EHR, electronic health record.

UMLS concepts was chosen to minimise false negatives in concept identification; relying on the user decisions about findings and the limited set of UMLS concepts to minimise false positives (table 2).

Creating simulated cases

Three cases of increasing complexity were created using real but deidentified clinical phenotypic and time course data from medical notes of Geisinger patients with known genetic diagnoses (online supplemental table 1). Cases were selected for conditions of varying complexity yet relatively common in the context of rare disease and

where diagnosis might be difficult using phenotype alone. Simulated cases were created by research assistants trained in capturing information from the EHR, supervised by a practicing Geisinger clinician certified in genetics and informatics. The three final cases were reviewed by a second Geisinger physician certified in genetics and informatics prior to user testing.

Case vignettes for the test scenarios assumed that some patient characterisation was previously noted by the clinician and genomic results were now available and could be interpreted with clinical information available in the

Table 2 Solutions for minimising false positives and negatives identified through NLP and DDSS by clinician review

False negative/positive problem	Solution included in GPACSS
Minimising false negatives on NLP flagging of findings	<ul style="list-style-type: none"> Include parent and child codes (eg, finding of intellectual disability in DDSS includes codes for developmental delay and particular types of intellectual disability).
Minimising false positives through the DDSS Usefulness metric	<ul style="list-style-type: none"> Use DDSS usefulness algorithm³⁰ to display flagged findings; thus prioritising data of greater relevance and de-prioritising data of low relevance for clinician review.
Minimising false positives through clinician verification	<ul style="list-style-type: none"> Use flag icon to indicate findings identified through NLP (figure 3). Clinician clicks the flag icon to display information needed to assess reliability, presence or absence, and onset. Information displayed from the EHR includes date of chart note, observer identity and three sentences of chart note (sentence with finding plus preceding and subsequent sentence).

DDSS, diagnostic decision support system; EHR, electronic health record; GPACSS, Genotype-Phenotype Archiving and Communication System with SimulConsult; NLP, natural language processing.

EHR (online supplemental figure A). For the three cases, a total of five findings were used as initial information before the genomic results, with three (one per case) being flagged findings identified through NLP. This created a 'near live'²⁴ experience within the simulated EHR for user testing while limiting the expense and time of EHR integration during this preliminary phase.

User testing methods

Participants

GPACSS is both a DDSS and communication tool to facilitate utilisation of genomic and phenotypic information available in the EHR by all clinicians to improve patient care within a healthcare system. Therefore, we purposively selected primary testers from Geisinger staff representative of current end users of the genome-phenome analyzer. Because a limited number of individuals at Geisinger regularly engage in using genomic information for differential diagnosis, we followed guidance recommending 3–5 evaluators for preliminary usability testing.²⁵ A group of secondary testers (inclusive of a pilot tester) with other roles in the genetic testing and interpretation process were purposively selected for potential broader utilisation in the healthcare system.

Testing sessions

At the beginning of each session, testers viewed a 4 min training video (<https://simulconsult.com/videogpacss>) beginning from saved patient findings, then importing a VCF, and review of flagged findings to make a diagnosis and create a customisable patient-friendly RoR report.

A semistructured interview guide (online supplemental file 3) was created to elucidate factors relevant to human factors in CDS design (information, interaction, interface)^{1 5 26} and organisational implementation (acceptability, perceived need, feasibility, workflow fit).²⁷ We used a think aloud²⁴ approach where testers were asked to verbalise thoughts while using the GPACSS prototype with the interviewer asking questions as needed and at key points in the testing to create a cognitive walkthrough with heuristic evaluation.^{25 28} Testers were invited via direct contact from study staff and provided a description of the study. At the beginning of each session, study staff reviewed a study information sheet and obtained verbal consent to participate. Test sessions lasted 2 hours and testers received a US\$100 gift card.

An experienced interviewer (AKR) and observer (MAW) from Geisinger worked with each tester to imagine using GPACSS for each test scenario. The interview and process were piloted with a cancer genetic counsellor reviewing one test vignette. At the end of the session, testers were asked a series of study-specific questions using a 0–10 rating scale (hard to easy) to rate the overall usefulness, satisfaction, and navigation. Transcripts were created from the audio portion of each session and the computer screen was video recorded to capture tester movement through GPACSS.

Analysis

Two Geisinger coders (MAW and JCR) viewed each user test session recording, read transcripts and created a codebook of themes identified across sessions. Transcripts were coded and the corresponding quotes were organised into a matrix using the three categories of CDS components (information, interface and interaction) identified by Miller *et al.*¹ and categories of acceptability, perceived need, feasibility and workflow fit according to Rogers' Diffusion of Innovations in organisations constructs.²⁷ Coders analysed transcripts independently and reviewed for agreement with discrepancies resolved by the primary author.

RESULTS

Three clinicians currently using genomic information to diagnose patients participated as primary testers: a paediatric geneticist (orders exomes daily), internal medicine physician (orders 4–5 exomes per month) and a paediatric genetic counsellor. Three additional clinicians participated as secondary testers; representing broader usability within the healthcare system: the pilot tester (cancer genetic counsellor), a laboratory director (conducts variant interpretation) and a laboratory genetic counsellor (conducts variant analysis).

GPACSS usability: human factors of CDS design

Overall impression of the prototype was positive. Testers raised general issues relevant to human factors in CDS design.¹⁵

Interface

Testers liked the flagged findings (figure 3), the contextual information for each mention in the EHR, and the rank ordering of flagged findings by usefulness. The visualisation of the evolving differential diagnosis and the automated RoR report for sharing with patients and referring clinicians, including the ability to save and access this report from the EHR were also appreciated.

The interface was noted to be complex, but testers stated this was expected due to the inherent complexity of genetic diagnosis and that they anticipated a learning curve to develop proficiency. Placement, positioning and the multiple presentation layers (text and graphics in the interface)¹ were well liked. In particular, the 'Assess diagnosis' display was noted as valuable because it made transparent the logic used by the DDSS in comparing patient findings to information about the disease. Of note, each tester interpreted differently the meaning of the graphical bars and shading, however, this did not hinder their ability to make the diagnosis, and the bar itself was appreciated as a design feature. To help with interpretation, more labelling was suggested (table 3).

Interaction

Testers were thoughtful and purposeful using GPACSS. Notably, in case 3 (the most complex case), one primary

Table 3 GPACSS usability: human factors of CDS design and Organisational implementation factors through tester Experiences*

Human factors of CDS design	Interface	<p>'More training would be good... unless I was doing it all the time for all of my patients, every step, I might not realise that some of the features are available...' (Tester 3)</p> <p>'These bars are different lengths, so I assume it's having something to do with frequencies... so I'm not sure why this part is purple...if there were something [on the assess diagnosis tab] that said this is 100% over here and this is 0% over here, that would kind of help, if I knew that that was the case... I'm not sure what these other colors are referring to.' (Tester 5)</p>
	Interaction	<p>'To me, the green bar in it shows me they are confident that this genetic variant aligns with the phenotypic markers that we have identified. I don't necessarily know how far the bars will tell me they're confidence in pathogenic vs VUS.' (Tester 3)</p> <p>'It's going to take a lot to learn. A lot of clicking back and forth and it's not super intuitive but I get it. So, the report gets generated and that becomes part of the record. I can see how that can be helpful because it has now particular phenotypical diagnosis and even genetic finding' (Tester 2)</p>
	Information	<p>'The term pertinent gene zygosity is not something I would normally make part of my lexicon...I have a general sense of the term zygosity but I can't remember the last ten years using that term in any of my discussions in clinical care or genetics in some of the cases I found what's their zygosity' (Tester 2)</p> <p>'But the variant severity score doesn't mean anything to me personally. To me it's easier to know, if you verify know the true classification they are giving it... Pathogenic, likely pathogenic VUS benign.' (Tester 3)</p> <p>'I think this one is nice [the 'Mention' displayed in a flagged finding]. Whenever someone says it had been noticed earlier by, it's nice when someone is talking about their niece or nephew, or like a proband cousin, they are saying they had myopia and I remember them having glasses before they were 5 years old.' (Tester 3)</p>
Organisational Implementation Factors	Acceptability	<p>'I would use it most of the time. To me, this is the frontier of genomic medicine and I look at my role as not only taking care of a patient but figuring out how we make genomics part of everyday medical practice. The useful things in the chart, genetics people can now get to right away'. (Tester 2)</p> <p>'...Typing them up, writing the summary [of all the patient findings in the chart]. If I could see what's been flagged in the chart, see what has not actively been flagged and decide do I need to go back and look at it or not. It would save my time' (Tester 3)</p> <p>'I think the interface is really good, in that you have that ability to explore those variants that may or may not make it on the reports that we get now, so you can drill deeper if you want. (Tester 5)</p>
	Perceived Need	<p>'...The report is a great idea for highlighting why you think it's [the care instructions] important, [in] a standard format... The average primary care physician that gets the genetic testing reports, says I don't know what this means at all. I think this [the Prognosis Table] is a step towards making it more understandable.' (Tester 2)</p> <p>'Everything's there [in the chart] and the question is how easy is it to find. I'm sure if you're a malpractice lawyer you get very good at pulling stuff out of these charts and asking why didn't you see that. Yet I can't look at everything.' (Tester 2)</p> <p>'This is stuff that you are doing anyway... you could make your note a lot shorter and just refer to that document [the automated Summary] ... I like the idea that you can explore. Clinical genetics now is limited on time.' (Tester 5)</p>
	Workflow Fit	<p>'It's nice because it helps guide me... it's very easier for me to realize that Prader-Willi is associated with narcolepsy...' (Tester 3)</p> <p>'I think the nice part is I don't have to go searching myself to find all the signs and symptoms associated with it and potentially miss something, that I may not know is a less common finding or feature of the condition. That actually could be beneficial for a provider or for us to give to the testing lab, to say these are all symptoms that we see, and then analyzing the data' (Tester 3)</p>

*Comments from primary user-testers only (testers experienced with differential diagnosis of genetic conditions through sequencing): n=3; paediatric genetic counsellor, paediatric geneticist, internist ordering 4–5 exomes in the past month.
CDS, clinical decision support; GPACSS, Genotype-Phenotype Archiving and Communication System with SimulConsult.

tester did not immediately choose the top diagnosis offered by GPACSS. Supported by the data displayed, the tester indicated that to make a definitive diagnosis they would next evaluate for the second-ranked disease—as that condition had a test that was easy and accurate and the condition was also more treatable—indicating utilisation of the DDSS as intended and consistent with clinical diagnostic decision making.

Testers initially expressed concern around ‘too many clicks’ and ‘click fatigue’ but noted as they progressed through the cases that the clicking was unavoidable and necessary. For example, they saw value in taking the time to correctly specify onset information (which requires clicking and cognitive load in the DDSS), as this is part of the genetic diagnostic process. ‘Cognitive Load’ in DDSS testing refers to additional thinking required to interact with the tool, and the general recommendation is to minimise this in CDS design.¹ Testers who commented on the cognitive load required to review flagged findings and choose age of onset noted the cognitive load as similar to completing this task without GPACSS.

Information

Testers appreciated resources such as the hover feature that revealed synonyms to findings and requested even more hovers and infobuttons. Confusion over some terminology occurred, notably ‘zygosity’ and ‘severity score,’ when reviewing the genomic variants; as only some testers located the explanatory resource for these terms.

The fact that the EHR ‘Mentions’ displayed in flagged findings were sometimes triggered by parent or by child concepts was noticed by all testers, and some stated the findings used in the DDSS were not as granular as they were expecting. Regardless, testers recognised and emphasised the importance of being able to review the ‘Mention’ information from the EHR and manually adjust for any false positives and false negatives from the NLP process.

GPACSS usability: organisational implementation factors

Acceptability

For the primary testers, satisfaction averaged 8.5 out of 10 (range 8–9.5) and navigation ease averaged 8 out of 10 (range 7.5–9). All three felt GPACSS would save time throughout the clinical process, with one primary tester estimating it at 30%–50%. Specific value in time saved was noted for chart review by all testers.

Perceived need

The RoR report and detailed prognosis table²⁰ generated in each scenario was highly valued for being standardised and for its ability to communicate complex genetic information to patients and other clinicians (table 3). The RoR report was also noted as an improvement over current laboratory reports; with one tester stating it was ‘where the most utility would be’ (Tester 4).

Testers exhibited learning and familiarity with GPACSS as they progressed through the testing session;

appreciating the DDSS assistance as each vignette increased in complexity; noting ‘It takes it [clinical diagnosis and diagnostic thinking] to a higher level’. [Tester 2]. Primary testers expressed readiness to adopt the tool in clinical practice; and one (paediatric geneticist) suggested GPACSS could also serve as a differential diagnosis training tool for medical students and residents in their clinic.

Two secondary testers (lab director and variant analyst) expressed enthusiasm that GPACSS could fill a need for in-house sequencing laboratories because full EHR data would be available during sequence interpretation. These testers also hypothesised that the ability to periodically re-analyse an existing VCF in minutes using GPACSS would improve the diagnosis rate over time.

Workflow fit

The three primary testers noted that the GPACSS process as tested fit with their clinical workflow diagnosing genetic conditions. As an added benefit, they described how using GPACSS also helped them learn about diseases and associated findings with which they were less familiar (table 3).

The three secondary testers questioned GPACSS fit with a clinical genetic testing workflow in which only a report with variants labelled as to pathogenicity and association with a condition (implying a clinical diagnosis) is received from an external lab. However, they did identify value and possible workflow fit for situations with uncertainty as to the diagnosis after sequencing or where flagged findings and the usefulness ranking would allow clinicians to review the EHR with flagged findings in light of the genomic information to make the diagnosis.

DISCUSSION

We provide preliminary evidence through user testing in a simulated real-world clinical workflow that the combination of NLP with a CDS tool optimised to support the clinical process of differential diagnosis may address the needs of those involved in this complex task. Such assessment of fit is critical if CDS is to fulfil the promise of standardising and improving care.¹⁴⁵⁸

Technical issues related to the interface and interaction of CDS design were minor and fixable; as were issues with design layout. Despite initial remarks on the number of clicks and cognitive load, testers acknowledged these as necessary to the genetic diagnosis process and no different than without the DDSS. Other issues related to terminology and usability could be solved and evaluated in future usability studies through a combination of training, added infobuttons and experience using GPACSS. Some of the technical gaps noted and additions requested by testers are addressed within GPACSS, however, the 4 min training video was created to provide enough instruction only to facilitate user testing. These results, therefore, provide direction for training and ongoing reference materials for future implementation.

For CDS to be acceptable and implemented by clinicians and organisations, it must fit with the real-world workflow and must present a solution to a perceived need.^{5 27} All primary testers identified ways GPACSS added such value and fit and noted ways GPACSS filled multiple needs in their diagnostic workflow. Workflow fit was highest among primary testers but opportunities for workflow fit were described by all testers. GPACSS was also noted as acceptable for implementation by all testers regardless of individual issues identified and suggestions for technical improvements.

LIMITATIONS

To facilitate user testing of GPACSS in the context of clinical workflow prior to full integration and implementation, simulations of the real-world were required. Because this study used the Logica EHR simulation, benefits or drawbacks of GPACSS in a production EHR could not be directly observed. Also, full annotations for the causal variants were not included in the variant table for the simulated patients limiting full assessment of the value of the DDSS in variant interpretation. This impacted the understanding of the 'severity score' by all testers, as the annotation information that would have been provided for a real patient was not included for the simulated cases. Finally, the generic cTAKES NLP using the UMLS concepts found only 20 of the 30 (67%) pertinent positive concepts within the test cases that a paediatric neurologist (MMS) identified manually. This was sufficient for GPACSS to generate the correct differential diagnosis for user testing, as further enrichment of the generic NLP to improve detection and avoid false positives was out of scope for this preliminary user testing.²⁹ Subsequent automated search for UMLS terms for flagging and addition of a separate stage of text search enrichment for terms missed by the NLP such as 'tall' improved NLP yield to 30 of 30 (100%).

This simulated EHR and user testing were a necessary first step and provide data to guide implementation of GPACSS. NLP improvements and additional beta testing within an actual EHR, in real-world clinical workflows, with real patient results and in real-world clinical workflows will be necessary to fully assess individual user-level and organisational-level facilitators and barriers to use, implementation and impact on clinical care. Such studies are currently in progress.

CONCLUSIONS

This study provides preliminary evidence for the usability, workflow fit, acceptability and implementation potential of a DDSS that includes machine-assisted chart review. Overall, responses suggest the GPACSS prototype is usable based on technical CDS and human-centred design criteria, addresses perceived clinical need, and has good fit within the real-world clinical workflow of genetic testing and diagnosis. Further development is needed to

improve usability for multiple clinical stakeholders and organisational implementation.

Author affiliations

¹Genomic Medicine Institute, Geisinger Health System, Danville, Pennsylvania, USA

²Intermountain Precision Genomics, Intermountain Healthcare, St. George, Utah, USA

³SimulConsult, Inc, Chestnut Hill, Massachusetts, USA

⁴University of Utah, Salt Lake City, Utah, USA

⁵Jackson Laboratory for Genomic Medicine, Farmington, Connecticut, USA

⁶University of Connecticut, Farmington, Connecticut, USA

Twitter Alanna Kulchak Rahm @Alanna_Rahm

Contributors All authors reviewed the final manuscript for submission and contributed to the study as follows: AKR contributed to the study design, conducted user testing sessions, and led the analysis, interpretation and writing of manuscript. NAW contributed to the study design, created the simulated clinical environment and the deidentified patient records, contributed to the NLP testing and adaptation of the DDSS and manuscript revisions. TNP was the lead developer on the NLP and the integration with the SimulConsult API. CJ, TJ, TNP and JP contributed to NLP testing and creation of the simulated clinical environment and manuscript revisions. MAW contributed to project management, user testing recruitment, qualitative data analysis and interpretation and review of manuscript. JCR contributed to project management, IRB applications, NLP testing and creation of the simulated clinical environment and manuscript writing. MSW contributed to study design, simulated clinical environment, NLP testing and adaptation of the DDSS, user testing design and recruitment, data analysis, interpretation and manuscript writing. LKF contributed to project management, study design and manuscript revisions. PNR assisted in the tagging of DDSS findings with HPO codes and created the new codes when needed. MMS designed and implemented the adaptation of the clinical decision support tool to interact with the NLP, designed and implemented the strategy for HPO and UMLS coding, did the manual chart review to identify findings that should have been recognised by NLP, and contributed to the study design, and manuscript revision.

Funding This study was supported by the National Human Genome Research Institute of the National Institutes of Health under the Small Business Innovation Research (SBIR) Program, Award Number 1R43HG010322-01 (principal investigator: MMS).

Disclaimer The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing interests LKF and MMS have an ownership stake in SimulConsult. MMS is PI of the NIH-funded Small Business Innovation Research (SBIR) grant. LKF and MMS provided the DDSS software to Geisinger for this research, including the modifications to enable the NLP and flagged findings, but were not involved in the user testing evaluation and analysis.

Patient consent for publication Not required.

Ethics approval The user testing protocol was reviewed and approved by the Geisinger IRB (#2018-0445).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Data include qualitative interview transcripts which are not permitted for reuse.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Alanna Kulchak Rahm <http://orcid.org/0000-0003-2513-9298>

REFERENCES

- 1 Miller K, Mosby D, Capan M, *et al.* Interface, information, interaction: a narrative review of design and functional requirements for clinical decision support. *J Am Med Inform Assoc* 2018;25:585–92.
- 2 Magrabi F, Ammenwerth E, McNair JB, *et al.* Artificial intelligence in clinical decision support: challenges for evaluating AI and practical implications. *Yearb Med Inform* 2019;28:128–34.
- 3 Bright TJ, Wong A, Dhurjati R, *et al.* Effect of clinical decision-support systems: a systematic review. *Ann Intern Med* 2012;157:29–43.
- 4 Moxey A, Robertson J, Newby D, *et al.* Computerized clinical decision support for prescribing: provision does not guarantee uptake. *J Am Med Inform Assoc* 2010;17:25–33.
- 5 Miller K, Capan M, Weldon D, *et al.* The design of decisions: matching clinical decision support recommendations to Nielsen's design heuristics. *Int J Med Inform* 2018;117:19–25.
- 6 Moja L, Liberati EG, Galuppo L, *et al.* Barriers and facilitators to the uptake of computerized clinical decision support systems in specialty hospitals: protocol for a qualitative cross-sectional study. *Implement Sci* 2014;9:105.
- 7 Liberati EG, Ruggiero F, Galuppo L, *et al.* What hinders the uptake of computerized clinical decision support systems in hospitals? A qualitative study and framework for implementation. *Implement Sci* 2017;12:113.
- 8 Sutton RT, Pincok D, Baumgart DC, *et al.* An overview of clinical decision support systems: benefits, risks, and strategies for success. *NPJ Digit Med* 2020;3:17.
- 9 Williams MS. Perspectives on what is needed to implement genomic medicine. *Mol Genet Genomic Med* 2015;3:155–9.
- 10 Shyr C, Kushniruk A, van Karnebeek CDM, *et al.* Dynamic software design for clinical exome and genome analyses: insights from bioinformaticians, clinical geneticists, and genetic counselors. *J Am Med Inform Assoc* 2016;23:257–68.
- 11 Segal MM, Williams MS, Gropman AL, *et al.* Evidence-Based decision support for neurological diagnosis reduces errors and unnecessary workup. *J Child Neurol* 2014;29:487–92.
- 12 Welch BM, Kawamoto K. Clinical decision support for genetically guided personalized medicine: a systematic review. *J Am Med Inform Assoc* 2013;20:388–400.
- 13 Grebe TA, Khushf G, Chen M, *et al.* The interface of genomic information with the electronic health record: a points to consider statement of the American College of medical genetics and genomics (ACMG). *Genet Med* 2020;22:1431–6.
- 14 Segal MM, George R, Waltman P, *et al.* Clinician-centric diagnosis of rare genetic diseases: performance of a gene pertinence metric in decision support for clinicians. *Orphanet J Rare Dis* 2020;15:191–91.
- 15 Segal MM, Abdellateef M, El-Hattab AW, *et al.* Clinical pertinence metric enables hypothesis-independent genome-phenome analysis for neurologic diagnosis. *J Child Neurol* 2015;30:881–8.
- 16 Issa MY, Chechacz Z, Stanley V, *et al.* Molecular diagnosis in recessive pediatric neurogenetic disease can help reduce disease recurrence in families. *BMC Med Genomics* 2020;13:68.
- 17 Williams MS, Williams JL, Rahm A. *Evaluating the impact of a user-friendly report of genetic testing results for parents of children with genetic disorders*. Washington, DC: Patient-Centered Outcomes Research Institute (PCORI), 2018.
- 18 Williams JL, Rahm AK, Zallen DT, *et al.* Impact of a Patient-Facing enhanced genomic results report to improve understanding, engagement, and communication. *J Genet Couns* 2018;27:358–69.
- 19 Williams JL, Rahm AK, Stuckey H, *et al.* Enhancing genomic laboratory reports: a qualitative analysis of provider review. *Am J Med Genet A* 2016;170A:1134–41.
- 20 Segal MM, Rahm AK, Hulse NC, *et al.* Experience with integrating diagnostic decision support software with electronic health records: benefits versus risks of information sharing. *EGEMS* 2017;5:23.
- 21 Kassirer JP. Teaching clinical medicine by iterative hypothesis testing. let's preach what we practice. *N Engl J Med* 1983;309:921–3.
- 22 Eddy DM, Clanton CH. The art of diagnosis: solving the clinicopathological exercise. *N Engl J Med* 1982;306:1263–8.
- 23 Savova GK, Masanz JJ, Ogren PV, *et al.* Mayo clinical text analysis and knowledge extraction system (cTAKES): architecture, component evaluation and applications. *J Am Med Inform Assoc* 2010;17:507–13.
- 24 Richardson S, Mishuris R, O'Connell A, *et al.* "Think aloud" and "Near live" usability testing of two complex clinical decision support tools. *Int J Med Inform* 2017;106:1–8.
- 25 Khajouei R, Zahiri Esfahani M, Jahani Y. Comparison of heuristic and cognitive walkthrough usability evaluation methods for evaluating health information systems. *J Am Med Inform Assoc* 2017;24:e55–60.
- 26 Nielsen J. Enhancing the explanatory power of usability heuristics. *Proc ACM CHI'94 Conf*, Boston, MA, 1994:152.
- 27 Rogers EM. *Innovation in organizations. diffusion of innovations*. 5th ed. New York, NY: Simon & Schuster, Inc, 2003: 402–35.
- 28 Khajouei R, Farahani F. A combination of two methods for evaluating the usability of a hospital information system. *BMC Med Inform Decis Mak* 2020;20:84.
- 29 Demner-Fushman D, Mork JG, Shooshan SE, *et al.* UMLS content views appropriate for Nlp processing of the biomedical literature vs. clinical text. *J Biomed Inform* 2010;43:587–94.
- 30 Segal MM. 2004 patent 6,754,655.