

COVID-19: are the elderly prepared for virtual healthcare?

Ahmed Ezzat,¹ Harpreet Sood,² Josephine Holt,¹ Hashim Ahmed,¹ Matthieu Komorowski ³

To cite: Ezzat A, Sood H, Holt J, *et al.* COVID-19: are the elderly prepared for virtual healthcare? *BMJ Health Care Inform* 2021;**28**:e100334. doi:10.1136/bmjhci-2021-100334

Accepted 02 March 2021

We follow with interest the unprecedented shift towards virtual healthcare during the COVID-19 pandemic. We echo concerns reported by colleagues of the fine balance between a need for global initiatives in cutting traditional red tape to enable rapid deployment of virtual health infrastructures versus the potential risks to quality of patient care that might occur when the patient is not physically in the same room as the clinician.¹

We have also highlighted the potential clinical, socioeconomic and environmental benefits of virtual consultations in secondary care.² However, with approximately one million consultations occurring each day in primary care in England alone, and 60% of primary care service users above the age of 60 years, there is a concern that elderly populations throughout the world will be disadvantaged in access to these virtual services due to lower proficiency with the tools of communication technology.^{3,4} For instance, the UK Office of National Statistics reports only 50% ownership of smartphones in those aged 55 and over versus up to 95% in the 16–24 age group.⁵

Besides the potential for accrued health risks from reduced physical interactions with healthcare services, we are concerned that older patients are at greater risk of social isolation and potential worsening mental health. If the sudden transition to remote consultations, rightly necessitated by a pandemic, is continued, then these patients may miss out on essential physical encounters including those crucial for chronic disease management. Indeed, for many patients who live alone, visits to health clinics constitute a social occasion to connect face to face with peers and carers.

So, when planning for a digital healthcare system beyond the immediacy of the COVID-19 pandemic, and which incorporates a greater role for remote consultations, careful consideration as to the impact to elderly and vulnerable populations is required. This is especially relevant during a second COVID-19 peak of infections and new local lockdown measures. Health policies can support this necessary technological revolution

in this age group with special consideration to their premorbid conditions, such as arthritis, visual impairment, or cognitive impairment. If planned well and delivered robustly, this virtual shift could also represent a fortuitous opportunity to widen access for the elderly in both ownership and use of information and communication technology. Access to virtual consultations, in the ageing population, emphasises the quality of life benefits, fostered through the adoption of technology. It is part of a growing trend of elderly people who not only receive healthcare but also stay in touch with friends and family at distance and reduce isolation.

Twitter Matthieu Komorowski @matkomorowski

Contributors All authors contributed equally.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Matthieu Komorowski <http://orcid.org/0000-0003-0559-5747>

REFERENCES

- 1 Webster P. Virtual health care in the era of COVID-19. *Lancet* 2020;395:1180–1.
- 2 Edison MA, Connor MJ, Miah S. Understanding Virtual Urology Clinics: A Systematic Review [published online ahead of print, 2020 May 28]. *BJU Int* 2020.
- 3 Available: <https://www.ethnicity-facts-figures.service.gov.uk/uk-population-by-ethnicity/demographics/age-groups/latest> [Accessed on 03/06/2020].
- 4 Available: <https://www.england.nhs.uk/five-year-forward-view/next-steps-on-the-nhs-five-year-forward-view/primary-care/> [Accessed on 03/06/2020].
- 5 Available: https://www.ofcom.org.uk/_data/assets/pdf_file/0022/117256/CMR-2018-narrative-report.pdf [Accessed on 17/10/2020].



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Surgery and Cancer, Faculty of Medicine, Imperial College London, London, UK

²Health Education England, London, UK

³Intensive Care Unit, Charing Cross Hospital, Imperial College Healthcare NHS Trust, London, UK

Correspondence to

Dr Ahmed Ezzat; a.ezzat@nhs.net