Information processing by community health nurses using mobile health (mHealth) tools for early identification of suicide and depression risks in Fiji Islands

Vimla Lodhia Patel,^{1,2} Mariel Halpern,¹ Vijayalakshmi Nagaraj,¹ Odille Chang,³ Sriram Iyengar,⁴ William May⁵

To cite: Patel VL, Halpern M, Nagaraj V, *et al.* Information processing by community health nurses using mobile health (mHealth) tools for early identification of suicide and depression risks in Fiji Islands. *BMJ Health Care Inform* 2021;**28**:e100342. doi:10.1136/bmjhci-2021-100342

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjhci-2021-100342>).

Received 28 June 2021
Accepted 25 October 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr Vimla Lodhia Patel;
vpatel@nyam.org

ABSTRACT

Objectives High rates of depression and suicide and a lack of trained psychiatrists have emerged as significant concerns in the low-income and middle-income countries (LMICs) such as the Pacific Island Countries (PICs). Readily available smartphones were leveraged with community health nurses (CHNs) in task-sharing for early identification of suicide and depression risks in Fiji Islands, the largest of PICs. This investigation examines how CHNs can efficiently and effectively process patient information about depression and suicide risk for making diagnostic and management decisions without compromising safety. The research is driven by the theoretical framework of text comprehension (knowledge representation and interpretation) and decision-making.

Methods Mobile health (mHealth) Application for Suicide Risk and Depression Assessment (ASRaDA) was designed to include culturally useful clinical guidelines for these disorders. A representative sample of 48 CHNs was recruited and presented with two clinical cases (depression and suicide) in a simulated setting under three conditions: No support, paper-based and mobile-based culturally valid guideline support. Data were collected as the nurses read through the scenarios, 'thinking aloud', before summarising, diagnoses and follow-up recommendations. Transcribed audiotapes were analysed using formal qualitative discourse analysis methods for diagnostic accuracy, comprehension of clinical problems and reasoning patterns.

Results Using guidelines on ASRaDA, the CHNs took less time to process patient information with more accurate diagnostic and therapeutic decisions for depression and suicide risk than with paper-based or no guideline conditions. A change in reasoning pattern for nurses' information processing was observed with decision support.

Discussion

Although these results are shown in a mental health setting in Fiji, there are reasons to believe they are generalisable beyond mental health and other lower-to-middle income countries.

Conclusions Culturally appropriate clinical guidelines on mHealth supports efficient information processing for quick and accurate decisions and a positive shift in

Summary

What is already known?

- Smart mobile phones have supported healthcare applications by health care providers in lower to middle-income countries.
- The use of smart mobile phones can support automated calculations to reduce the total number of errors.
- Although there have been experiments in mental health settings, their impact has not been rigorously evaluated.

What does this paper add?

- The information processing involves filtering irrelevant information from the relevant, contributing to a lower cognitive load on the providers, thereby making the process efficient.
- Safety may be compromised at the expense of efficiency due to conceptual errors even though the total number of errors is reduced.
- The use of culturally appropriate guideline support on mobile phones can result in positive behavior change.
- Although these results are shown in mental health settings, there are reasons to believe they are generalised beyond mental health.

reasoning behaviour by the nurses. However, translating complex qualitative patient information into quantitative scores could generate conceptual errors. These results are valid in simulated conditions.

INTRODUCTION

Mental health (MH) disorders are globally recognised as a significant public health concern yet receive inadequate attention, and more than 80% of people experiencing mental illness live in low-income and middle-income countries (LMICs).¹

Pacific Island Countries (PICs) are classified as LMIC, where in recent years, depression and suicide have been reported to have reached an alarming level.² With fewer than 20 qualified psychiatrists serving more than 11 million people, this situation is only likely to worsen with pandemic and post-pandemic-related conditions. The PICs alone account for 25% of the global burden of disease: Death by suicide comprises approximately 25% of the total disease burden,³ and depressive disorders approximately 5.7%.¹

To increase coverage and access to evidence-based mental healthcare, community health nurses (CHNs), the front-line healthcare workers, assume a significant role in the early identification of depression and suicide risk in a collaborative care approach called 'task sharing'. Despite the relative success of task-sharing initiatives globally, there is a lack of consistency in the quality of MH care, where individuals who have severe mental illness are not appropriately diagnosed and treated by community health workers.^{4,5} In the community clinics in the PICs, outpatients come with different ailments, and the CHNs are challenged to identify the ones with mental health risks. This task generates a high cognitive load on these nurses, necessitating a need to leverage cheap and easily available technology, as support to offload some of the burden.

Given the widespread accessibility to smartphones in Fiji, it is appropriate to use them to support CHNs in MH care delivery by developing mobile-based screening applications to quickly screen and provide care to patients at risk. However, while mobile technology intervention have great potential to help address current problems, little is known about its added value to provide efficient, effective and safe care in the PICs.

The optimal design of mobile health (mHealth) interventions requires an understanding of the cognitive underpinnings of the use of any technological devices since the interventions are most often ad hoc and lack a sound empirical basis for investigations.⁶ The use of mHealth has been successfully undertaken to support nurses in providing bedside documentation and to assist in their usual workflow.^{7,8} Although, smartphones and related mobile technologies are recognised as flexible and powerful tools when used sensibly in augmenting human cognition, there is also a growing concern that continued use of these devices, may have a negative impact on users' cognition, namely, their ability to think and pay attention to specific tasks.⁹

Therefore, our studies are driven by theories and methods from cognitive science related with special focus on text comprehension (knowledge representation and interpretation) and patient management decisions within the cultural context.^{10,11} The cultural context refers to the social behaviour and norms practiced in the Fiji culture, including the peoples' knowledge, beliefs, language and customs. It is the way they think, understand and communicate, which is important to consider for building any successful interventional tools.

Qualitative–cognitive studies involves collecting and analysing data from a fewer number of representative participants, with a focus on their knowledge structures and use of cognitive strategies in processing patient information.^{12,13} Cognitive strategies are the specific methods that people use while solving problems. This process of eliciting underlying thoughts while reading or solving a problem requires a well-established procedure in psychological literature of collecting lengthy 'think-aloud' protocols from the nurses, as they interpret, identify and resolve the patient problem.¹⁴

This 'think-aloud' method is used extensively in psychological and educational research and offers unique approach to capture participant thought processes when interacting with technology and interventions.^{15,16}

METHODOLOGY

Rationale

We are using smaller sample sizes of participants with in-depth analysis in our studies rather than a large-scale epidemiological study, limiting our ability to run statistical analyses on the data. We aim to develop a basic understanding of a complex phenomenon under consideration before testing generalisable inferences from this data. It should be noted that in any complex clinical situation filled with idiosyncrasies, varied situations and unique constraints, there are a remarkable number of invariants on which we can capitalise once we identify the over-bidding phenomenon that is task and people invariant. Although our sample size is small, we attempt to get data saturation with qualitative methods.¹⁷

To characterise cognitive strategies underlying diagnosis for depression and suicide by CHNs under three conditions: no guidelines (current practices) as the base-line and two guidelines, paper and mHealth tools. Representation of patient information is different in the three conditions, and thus the information will be interpreted differently.¹⁸ The patient data in the paper-based support tool is in a narrative form. The data in the mobile phone is discrete, requiring a shorter time to process information and require different nature of inferences to be drawn for making diagnostic decisions.

We expect that the nurses using mHealth will (a) process clinical information in a shorter time and (b) generate fewer errors in the calculation for more accurate diagnostic and therapeutic decisions.

Study design and procedures

Using an experimental design, CHNs were assigned to one of three conditions, nurses using to guidelines on mobile phones, on paper, and the last group were given no guideline support (control, reflecting the current status).

In all conditions, CHNs were presented with two clinical case scenarios, one representing severe depression and the other representing a high risk for suicide (described in materials and measures below). The CHNs

	Clinical task	Experimental condition		Control condition
		Mobile-based guideline (N = 16)	Paper-based guideline (N = 16)	No guideline control (N = 16)
Clinical case Severe depression	1 Think aloud	✓	✓	✓
	2 Summary problem	✓	✓	✓
	3 Final Diagnosis	✓	✓	✓
	4 Recommendation(s) for treatment	✓	✓	✓
High risk for Suicide	1 Think aloud	✓	✓	✓
	2 Summary problem	✓	✓	✓
	3 Final diagnosis	✓	✓	✓
	4 Recommendation(s) for treatment	✓	✓	✓
Severe depression with guideline	1 Think aloud		✓	
	2 Summary problem		✓	
	3 Final Diagnosis		✓	
	4 Recommendation(s) for treatment		✓	
High risk for suicide with guideline	1 Think aloud	✓		
	2 Summary problem	✓		
	3 Final Diagnosis	✓		
	4 Recommendations for treatment	✓		

Figure 1 Study design and procedures for severe depression and suicide risks.

in the control condition were tasked with four clinical problem-solving activities: (1) Think aloud while reading through each case (referred to as the ‘think-aloud’ task), (2) summarise each case (referred to as the ‘summary problem’ task), (3) provide a final diagnosis (referred to as ‘final diagnosis’ task) and (4) provide recommendation(s) for treatment (referred to as ‘recommendation(s) for treatment’ task).

After completion of all four activities, only the CHNs in the two experimental conditions were given guidelines and asked to engage in the same four activities mentioned above using the respective guideline, as illustrated in figure 1. All nurses were required to read the clinical problem scenarios without any support from the mobile or paper-based guideline to get the base line data, before they were asked to use either mobile or paper-based technology.

Materials and measures

Clinical cases

Two clinical scenarios used in the study reflecting severe depression and suicide risk were developed by psychiatrists and primary care physicians based on what would be expected in real community clinics in Fiji. As an illustrated example, the case of severe depression is presented in figure 2.

Clinical guidelines

Pre-existing paper-based guidelines—the ‘Center for Epidemiological Studies Depression Scale’ (CES-D).¹⁹ for depression, and the ‘Suicide Behaviors Questionnaire-Revised’ (SBQ-R)²⁰ for suicide—were selected for the study. CES-D is a screening tool for depression developed by the CES and validated across various populations including Fiji.^{21 22} The guidelines are given in online supplemental appendices A and B. The questions (20-item for depression and four items for suicide risk) were converted for use by nurses (from self-evaluation) evaluated for cultural sensitivity and contextual appropriateness with the support of local clinicians, and updated via an iterative design process.

Ravi is 40-year-old Fijian male of Indian descent; employed as a bus driver; married with 3 children; living in Makoi.

PC: One-month duration of the following:

- Feeling worried,
- Insomnia and poor appetite,
- Headache,
- Easily fatigued
- Drinking kava and smoking suki more than usual

HPC: The above symptoms started after Ravi’s wife was diagnosed with breast cancer. She has been in and out of hospital for treatment, leaving Ravi to care for the children. He says he worries about the future a lot and whether his wife will recover. He feels guilty for not being more caring towards her and has frequent thoughts of death.

He finds it difficult to sleep at night and so drinks kava in the evening to help him sleep. Work has been hard for him as he is also having conflicts with his boss and finds that he has been smoking suki more than usual to help him cope with the stress. He finds he gets tired very easily and has frequent headaches. This has made it challenging for him to do his work and often misses work 1-2 times per week.

PMH/PAST PSYCHIATRIC HX: Nil

MEDICATIONS/ALLERGIES: Nil

SOCIAL/OCCUPATIONAL HX: Arranged marriage at age 22 years; wife is 2 years younger (housewife); have 3 children: aged 16 years (male); 14 years (male); 10 years (female); live with nuclear family in a settlement in Makoi (2-bedroom, tin-covered house with outdoor toilet and shower); achieved a Form 3 education; worked as a casual laborer immediately after leaving school. Employed as a bus driver for the past 18 years. Smokes suki daily 3-4 rolls for past 20 years; drinks alcohol on social occasions; Kava daily for the past 18 years; no marijuana use

ROS: Currently has difficulty falling asleep; has poor appetite (with loss of ~2 kg over the past month); Nil other problems.

MSE: Appears kempt; well groomed; Says he feels ‘okay’ but appears worried. Speech is relevant; coherent; but slow and low volume. He expresses concern for his wife’s health and claims he sometimes wishes ‘he could rest for good’ but doesn’t elaborate further. He is easily annoyed by questions asked. He denies having any perceptual disturbances and is cognitively intact. He has fair insight and judgment.

Note: HX History; PC presenting complaint; HPC history of presenting complaint; PMH past medical history; ROS review of symptoms; MSE mental status examination. Form 3 education = 8th Grade in Canada and US system; Suki = Native tobacco; Kava= A mildly intoxicated drink made from plant roots of the Pacific Islands.

Figure 2 Clinical case of severe depression in the Pacific Islands.

These guidelines were programmed into a mobile-phone application, *Application for Suicide Risk and Depression Assessment* (ASRaDA), and updated using a number of iterations with usability evaluation provided by CHNs. The left hand side of figure 3 shows the suicide guideline screen in the mobile phone, together with a nurse’s response.

The nurses read the questionnaire items aloud to a member of the research team who acted as the patient, and who provided a standard scripted response (typical of responses in Fiji) to the nurses. For example, the patient would respond to an item #5, ‘How often have you had trouble keeping your mind on what you were doing?’ with, ‘It has been really difficult to do my work or my usual chores’. This narrative response had to be translated into a quantitative number on a symptom frequency scale, from zero (rarely or none of the time; less than 1 day) to three (most or all of the time; 5–7 days). Total scores were added manually in the paper-based condition, but they were automatically calculated in the mobile condition.

Determining the criticality of guideline information

Two experts scored information in each of the two clinical cases as critical, important or additional supporting information for making the diagnoses. These items were then checked against the guideline items for alignment with the clinical aspects of the patient problem to capture

Figure 3 SBQ-R Screen in ASRaDA showing guideline questionnaire and nurse's response, and System Usability Scale.

the appropriate constructs, as intended. The criticality of guideline information is given in figure 4.

Mobile technology and development

The application development system, invented initially at NASA Johnson Space Center that enabled the development of media-rich training and clinical practice guideline applications without programming,^{22 23} was used to design and test a mobile phone based clinical guidance in rural health providers in India and in Colombia.^{18 24 25} In our current study, we leveraged concepts and experiences from this tested system to develop the suicide risk and depression assessment mHealth tool (ASRaDA) on an Android smartphone application, and loaded on Samsung J3 smartphones. Several versions of ASRaDA were tested in a controlled setting with a small cohort of CHNs, and refined iteratively into a final version, including responding to the 10-item System Usability Survey (SUS).^{26 27}

CHNs logged with the patient's name into the application, displaying guideline selection (see online supplemental appendix C, for an example of the mobile ASRaDA application guideline). Each participant was given an

identification number for privacy, where the mapping to the actual name was encrypted for security.

Participants

Forty-eight CHNs were recruited from an available pool of 5000 participants with the assistance of the Fiji Ministry of Health (MoH), according to the inclusion and exclusion criteria (race representation, relative fluency in English language, ages and years of experience). Since more women traditionally train as nurses, the possible recruitments were all women. The detailed eligibility criteria included: practicing or medically trained CHNs, who has satisfied requirements of Fiji's MoH, fluent in English (national language of Fiji), capable of using smartphone applications, between 25 and 55 years-old, and at least 5 years of nursing-related experience. Approximately 60% of the CHNs were local Fijians and 40% were Fijians of Indian origin, representing the per cent distribution of nurses in the community clinics. These percentages also align with statistics from prior research studies.^{28 29}

Only women traditionally train as nurses in the Pacific Islands. Representing the Fiji demographics, the recruits were approximately 65% i-Taukei (natives of Fiji) and 35% of Indo-Fijian origin. These percentages align with statistics from prior research studies.^{28 29} The community nurses had varied training and included nurses with general nursing (54%), mental health (23%) with WHO mhGAP³⁰ and other specialised training (79%) such as maternity or dermatology.

Methods of analysis

The data were collected and analysed using mixed methods approach that closely aligned with the tasks and the corresponding cognitive measures, as given in figure 5.

Audio recordings on the mobile phones were transcribed using 'Express Scribe Transcription Software' V.6.10 from NCH³¹ automated software and checked for accuracy by the local experts. Using thematic qualitative approaches with inductive and deductive coding, as well as specific theme development,^{32 33} the generated 'think-aloud' protocols were analysed. Initial emerging

	Symptom	Question Numbers	Type
Severe depression risk	Sadness (Dysphoria)	2, 4, 6	Critical
	Loss of Interest Anhedonia)	8, 10	Critical
	Suicidal Ideation	14, 15	Critical
	Loss of Appetite	1, 18	Important
	Insomnia	5, 11, 19	Important
	Thinking/Concentration	3, 20	Important
	Guilt (Worthlessness)	9, 17	Important
	Tired (Fatigue)	7, 16	Additional
	Movement (Agitation)	12, 13	Additional
Suicide risk	Thought or attempt of committing suicide	1	Critical
	Past thought of committing suicide	2	Critical
	Confiding thought of committing suicide	3	Important
	Likelihood of committing suicide	4	Important

Figure 4 Expert classification of level of criticality of information in clinical text and corresponding guideline(s) for depression and suicide risks.

- Step 1: Evaluation of two patient scenarios
 - Subjects asked to read case while thinking aloud, unaided by clinical guideline
 - Summarize current status of patient: Conceptual understanding
 - Provide diagnosis and management plan: Effectiveness
 - Time on task automatically measured: Efficiency
 - Think aloud protocols recorded: Reasoning pattern
- Step 2: Evaluation of the same two patients while using guidelines for support
 - Same procedure as above, but with guidelines made available
 - Half of subjects use paper-based guidelines (requires calculating diagnostic scores): Errors in calculation
 - Half of subjects use interactive guidelines on mobile phone (scores automatically calculated): Effectiveness
 - Time on task automatically measured: Efficiency

Figure 5 Mixed methods: tasks and measures.

themes were used to code the rest of the transcripts to create descriptions of all emerging themes that accurately reflected the data. Nurses' reasoning patterns were identified using method in previous research on guideline-assisted reasoning.^{12 13} The response protocol segmentation processes identified both the concepts used by the nurses and the relations among these concepts in their reasoning. Semantic networks were then constructed, where concepts and their relations are organised into graphical structures that represent the steps in their reasoning. These semantic networks convey two types of information: conceptual information (ie, the concepts used to solve a problem) and structural information (ie, how the concepts are related to each other) about how the subjects constructed their diagnoses and recommendations

The diagnostic accuracy coding was adapted from Patel and Groen³⁴ and comprised three categories: Correct (ie, completely accurate), partially correct (ie, some accurate diagnostic components) or incorrect (ie, not accurate).

These categories were evaluated as a function of the levels of nurses' MH training.

The summarisation protocols were analysed using formal methods of natural language representation, more specifically, concept and semantic analysis.³⁴ Next, the expert template for information criticality was used to code CHNs summary protocols. An illustrated example of the coding of these three summary narratives is given on figure 6.

An example of the first step in clinical text segmentation and coding of CHNs protocols for generated inferences using the depression scenario is given in online supplemental appendix D.

The times for task completion were automatically calculated on the smart phone recorders.

RESULTS

Time on task: task efficiency

Task efficiency is the time required to complete the task. The benchmark was the time on task with no guideline support (mobile or paper).

Figure 7 depicts time for task completion with paper and mobile-based guidelines for depression and suicide risks scenarios.

The average times for completion of the depression and suicide risk problems were the shortest when using the mobile technology, followed by the paper-based and the no guideline support condition. The use of mHealth-based guidelines was most efficient in supporting quick identification of suicide and depression risk patients.

Time on task: process efficiency

Process efficiency is the processing of patient information for depression and suicide risks as identified by concept density and accuracy of patients description in the problem summaries provided by the nurses.

	Summary	Evaluation	# of Concepts	Limitation	Recommendation	Advice	
No Guidelines	Staff Tavaitea doing afternoon shift. I have a case to hand over to you / Uuhh... Ravi is a 40 year old... uuhh Fijian male / Employed as a <u>bus driver with 3 kids</u> / They lived in Makou and uuhh... the according to Ravi even Ravi has been <u>feeling worried</u> and uuhh... poor appetite/ uuhh... <u>easily fatigued</u> and <u>drinking kava</u> and <u>suki</u> more than usual/ Uuhh... also the what has, has caused Ravi to change his behavior in terms of in terms of his work? He has been <u>missing work</u> for 1 to 2 week/ 2 times per week (and he also <u>facing conflict with his boss</u> / And he eats / uuhh... drinking kava more than usual/ Plus due to that, due to I think what causes that? is the wife is uuhh... <u>his wife being diagnosed with breast cancer</u> / plus his job / he was just a bus driver/ and he has 3 months to feed which is the 3 kids that he still looks after/ and they still are schooling/ and uuhh... He has been smoking <u>suki</u> /	Longer summary with both relevant and irrelevant information	28			"For Ravi, he should go for.... we have uh... counseling uh... counseling net, counseling service. Will offer Ravi a session just to show Ravi that that's not the end of it and also bringing up to the state that is required."	Counseling (to be able to cope with this situation)
Paper-based guidelines	"Ok, uuhm... Ravi, forty year old Fijian male/ uuhm... came in with ahh... symptoms of <u>feeling worried</u> , / uuhm... <u>insomnia</u> and <u>poor appetite</u> / <u>headache</u> / <u>easily fatigued</u> and his been <u>drinking kava and smoking suki</u> more than usual for the past one month / Uuhm... he lives with his wife/ who is uuhm... sick. / <u>lately diagnosed with breast cancer</u> / And 3 children, / all schooling / One 16 years, 14 years and 10 years / So he is the <u>only bread winner</u> in the house / He worked as a <u>bus driver</u> / and they live in a, they live by themselves/ so his the only that has been looking after the family financially. / And the wife is very sick / so he came with ahh... all his problems/ so his ahh... <u>really under stress</u> /"	Shorter summary with mostly relevant information	20	Guideline constrains the search space	"For Number 1, Severe score. Severe score, ahh I will refer patients refer patients for relevant Stress Management Ward at Suva and make sure he gets proper treatment at St. Gils."	Exactly as recommended, based on the score	
With Mobile Guidelines	"Ok uhm... to my oncoming staff I would tell her about this case/ like Mr. Ravi is a 40-year-old/ Fijian male of Indian descent/ He is employed as a <u>bus driver</u> / married with 3 children / and living in Makou / Uhm... for one month he has been <u>feeling worried</u> / his had <u>insomnia</u> and <u>poor appetite</u> / uuhh... <u>headache</u> / He is <u>easily fatigued</u> / and has been <u>drinking kava</u> and <u>smoking suki</u> / more than usual / Uhm... he presented to the health facility / and... according to him the all this problem started / when his <u>wife was diagnosed with breast cancer</u> /"	Shorter summary with mostly relevant and missing information	16	Mobile Guideline constrains the search space.	"Yes uh... refer patients to the relevant stress management ward / at Suva / or possibly St Gils, Lautoka Hospital / that's in Western Division / or Labasa Hospital in the Northern Division / In more isolated locations / make direct referral to the relevant stress management ward / for St. Gils / Patients will most likely commence on anti-depressants / and probably admitted /"	Exactly as recommended (probably based on the score)	

Figure 6 An example of analysis of summaries and recommendations as a function of guideline use by a community health nurses.

Figure 7 Time for task completion with paper and mobile based guidelines for depression and suicide risk. CHNs, community health nurses.

Summarisation tasks were analysed for knowledge organisation among the control and the experimental groups. There were fewer concepts identified in the summaries provided by CHNs in the guideline experimental conditions for depression (paper, $M=6.44$ (6.34); mobile, $M=11.11$ (3.52)) and suicide (paper, $M=3.69$ (7.04); mobile, $M=3.10$ (1.90)) problems than in the summaries under control condition (depression: $M=15.94$ (7.85) and suicide: $M=12.69$ (5.79)) for both depression and suicide risk cases (see [figure 8](#)). Most of the information in summaries with guideline support were important and critical only, whereas in the control condition, irrelevant information was also included.

Reasoning about the patient problems

Knowledge organisation drives the development of reasoning strategies for diagnoses. Thus, it is expected that the CHNs will show different reasoning patterns in

Figure 8 Mean number of concepts in summarization task using paper, mobile, and no guideline conditions. CES-D, Center for Epidemiological Studies Depression Scale; SBQ-R, Suicide Behaviors Questionnaire-Revised.

the control and in both the experimental conditions. As reported earlier, the transcripts were analysed using a formal method of natural language analysis, the semantic networks.¹³

Figure 9A presents an illustrated example of a community nurse's think-aloud protocol (nurse #1 with MH training) in the control group, working with the depression case. The figure shows a semantic network structure, where nodes are patient information given in the clinical text. The links are the directionality of inferences generated during the think-aloud process.¹⁸ She identified the key concepts associated with the scenario but attributed the financial crisis to the patient's 'bad' behaviour, which led to depression, and she recommended that the problem may go away if the patient stopped spending money on drinking kava and smoking suki.

Figure 9B presents an illustrated example of a think-aloud protocol of another community nurse (nurse #2 with MH training) with an equal level of training in MH when primed by the depression guideline. The nurse first considered severe depression, a significant factor as a possible diagnosis, and looked for confirming evidence in the patient scenario. Drinking more kava and smoking suki was considered a result of depression rather than the cause. She recommended that Ravi seek professional help.

With the use of paper-based guidelines, a similar pattern was found. This shift in reasoning was identified in 35% and 30% of the nurses working with depression scenarios in mobile and paper-based support, respectively, showing a specific positive trend in behaviour change. These patients were recommended for illness management according to the guidelines, which otherwise would have been missed. The rest of the nurses either did not generate rich enough data to conduct the analysis of natural language (NLP), as with the rest of the data, or did not show any change in their reasoning patterns. The suicide risk scenario guideline was too short, resulting in sparse think-aloud data, and was not amenable to formal analysis for reasoning patterns.

Errors of calculation

Errors were found in manually adding the total number of scores with the paper-based guideline for depression (correct score: 36.55, CHNs: 30.00 ± 2.6), and suicide risk (correct score: 11.15, CHNs: 12.15 ± 1.3). In this particular situation, the errors did not affect the recommended treatment modalities since a wide range of the scores for each treatment modality resulted in the same treatment recommendations. However, in some other situations, where the score range relates for treatment recommendation is much smaller, these errors will be of concern. There were no calculation errors when using mobile-based guidelines since the system automatically made the calculations and suggested the appropriate recommendations based on the total scores.

Figure 9 (A) Patient problem explanation prior to reading the mHealth guideline for depression. Key: rectangles represent concepts identified and the links represent connections between the concepts as inferred by the community health nurses. The arrows represent the directionality of reasoning. (B): patient problem explanation AFTER reading the mHealth depression guideline.

Diagnostic accuracy

Table 1 presents CHNs final diagnostic decision patterns after going through the patient problems.

In the case of severe depression, the majority of CHNs provided a partially correct final diagnosis. The four nurses, who provided the correct diagnosis, had recently received or were currently undergoing training in MH (WHO-based mhGAP programme).³⁰ However, two of these four nurses provided only partially correct recommendations for treatment. Five CHNs who provided incorrect diagnoses had not received MH training. Training in mental health programme appears to play an important role in diagnostic accuracy for identification of depression and suicide risk.

Errors of translation: From qualitative to quantitative information.

During our analysis, we discovered other form of errors when coding nurses' responses to the guideline questions (qualitative) into quantitative scales. These errors were generated during cognitive processing of patient information for translating them (interpreting) into paper and

mobile-based quantitative scales to determine the treatment recommendations.

Not all the nurses provided detailed 'think-aloud' data when computing this translation. Six nurses (total: 9), using paper-based guidelines and five (total: 7) using mobile-based depression guidelines, generated three to four errors in translation. An illustrated example of two such errors is given in table 2.

Suicide risk guideline was too short to capture such errors. Limitations in data collection restricted our ability to provide a more comprehensive information on these errors, except that such errors are pervasive due to the complexity of patient responses, which do not fit neatly into a set of quantities.

Usability

Sixteen nurses who used the ASRaDA application, 7 participants for CES-D and 14 for SBQ-R completed the questionnaire. From a maximum score of 100 on SUS usability scale, ASRaDA was scored at 85.36. The details of the usability study and its limitations are reported elsewhere.^{29 35} The left hand side of figure 3 shows the suicide guideline screen on the mobile phone, together with a nurses' response to the usability test.

DISCUSSION

Clinical guideline acts as a primer to help focus on fewer and more critical/important information, filtering out most irrelevant information.^{34 36} It constrains a broad memory search, making it easier and quicker to get to the target, due to reduced memory load. The ability to separate relevant information from irrelevant information is recognised as a hallmark of developing expertise, where the guidelines provide such support.

The conceptual shift in reasoning assists the nurses to narrow the relevant information search space and, quickly arrive at the diagnosis.³⁷ This change in the directionality of reasoning suggests a critical behaviour change from blaming the patient to attributing the problem to the nature of the illness. This pattern of behaviour has also been reported in other studies.⁵

Table 1 Final diagnosis provided by the community health nurses

	Clinical problems		
	Mental health (MH) Training level	Severe depression N (%)	High risk for suicide N (%)
Correct	MH training	4 (100)	0 (0)
	No MH training	0 (0)	0 (0)
	Total	4 (8)	0 (0)
Partially correct	MH training	8 (26)	12 (40)
	No MH training	22 (73)	18 (60)
	Total	30 (63)	30 (63)
Incorrect	MH training	3 (60)	0 (0)
	No MH training	2 (40)	7 (100)
	Total	5 (13)	7 (15)
Missing		9 (24)	11 (23)
Total		48	48

Table 2 Errors in translating narrative responses to the questions into quantity

Example 1: Nurse #07	Example: Nurse #08
MHb07: "Do you enjoy life?" Patient (Ravi): "Not at all." MHb07: "Most of the time. <u>Three</u> ." Interviewer: "So are you saying that most of the time he enjoys life, based on Ravi's answer?" MHb07: "Ok let me ask you the question again." "Do you enjoy life?" Interviewer: "Ravi says not at all." MHb07: "That means never so that's <u>zero</u> ."	MHb08: "Do you feel you are just as good as other people?" Patient (Ravi): "I have been feeling as unworthy as a criminal most of the time." MHb08: "Ohh...for that it will be <u>score 3</u> ." Interviewer: "Score 3?" MHb08: "Score 3." Interviewer: "So you are saying that Ravi feels just as good as other people most of the time based on his answer?" MHb08: "Oh no, no, he does not feel good at all so it will be score nil. <u>Zero</u> ."

The nature of patient information processing is different for nurses with and without mHealth-supported guidelines. The narrative nature of paper-based guidelines takes longer to process, but requires fewer inferences to be generated with fewer chances of any significant errors. The discrete nature of mobile-based guidelines helps focus on critical information faster. However, the user is required to generate more inferences in order to relate the discrete symptoms, to generating some form of patient narrative, resulting in a greater chance of generating errors of interpretation.³⁸

These results are shown to be true in our small sample of participants. With detailed analysis of fewer subjects, the results predict with some measure of confidence that community nurses with vastly different backgrounds, working in different settings, will generally experience similar problems when dealing with cognitive and technical performance pressures of varied characteristics. This detailed study enhances the reliability and rigour in generalising the study findings. The next step is to test them in the broader sample of nurses and implement and evaluate the tool in the clinical setting. The results are generalisable to other PICs (which share a relatively similar culture) and to other LMICs, as long as the tool is ecologically validated and tested for cultural sensitivity in the new context.

Impact of this research: Currently, treatment of those in PICs with mental health problems occurs when the disease is at an acute stage, for example, an unsuccessful suicide attempt, that necessitates hospitalisation. There are very few means or methods in the PICs to identify and treat at-risk persons at an early stage of the disease when treatment can be most effective. A significant benefit of our research is (a) increased task-sharing and empowerment of CHNs as the front-line care providers in mental health care and (b) likelihood of better mobile tool acceptability and sustainability in the clinical environment. The primary strength of the research is a novel application of cognitive, decision-making and safety aspects in mHealth design and user-centred testing process.

The targeted clinical problem, assessment of the risk of suicide in patients with depression, is an active public health issue everywhere in the world, with relevance to a variety of populations (for instance, war veterans and active military in the USA) and in virtually every context (outpatient or inpatient settings, primary care or rural environments).

CONCLUSIONS

The evidence from our exploratory study supports the hypothesis that culturally appropriate smartphone technology assists in the process of task-sharing with community health nurses, thereby reducing the burden on a limited number of available mental health professionals. The mHealth supports efficient and effective screening of patients with severe depression and suicide risk, with minor safety concerns. Furthermore, the results from the cognitive analysis show that culturally appropriate clinical guidelines for mental health risks trigger a positive trend in behaviour change.

LIMITATIONS AND CHALLENGES

There are always challenges when intervening in any complex and culturally diverse environment. This situation is exacerbated in resource-limited conditions such as in LMICs. One such challenge is about implementation in the clinical practices. When implementing ASRaDA into the community clinics in the PICs, these challenges will be salient, and will need to be carefully monitored for any unintended consequences.

One of our experimental tasks required the nurses to 'think-aloud' while reading the clinical cases. As in several other cultures, people do not often read aloud in a classroom in the PICs. This resulted in the nurses processing more information at the level of text syntax rather than the semantics, making it difficult to gauge more accurately, their understanding of the clinical problem. Training the nurses to think aloud while doing their tasks was not trivial, leading to some sparse and missing information during data collection, limiting our ability to collect the complete data set.

Author affiliations

¹Center for Cognitive Studies in Medicine and Public Health, New York Academy of Medicine, New York, New York, USA

²Biomedical Informatics, Columbia University Vagelos College of Physicians and Surgeons, New York, New York, USA

³Mental Health, Child and Adult Medicine, Fiji National University College of Medicine Nursing and Health Sciences - Tamavua Campus, Suva, Rewa, Fiji

⁴Internal Medicine, The University of Arizona College of Medicine Phoenix, Phoenix, Arizona, USA

⁵Dean's Office, Fiji National University College of Medicine Nursing and Health Sciences - Tamavua Campus, Suva, Rewa, Fiji

Twitter Vimla Lodhia Patel @VimlaLPatel

Contributors VLP conceptualised, set the direction of the investigation and drafted the final manuscript. MH and VN assisted in data analysis, creating figures and drafting of the primary manuscript. OC and WM provided the clinical support in monitoring the entire study at the clinical site. SI developed the mobile application and conducted the usability studies. VLP, the guarantor accepts full responsibility for the finished work and the conduct of the study, as well as the control the decision to publish.

Funding This work was supported by the National Institute of Mental Health of the National Institutes of Health under Award Number R21MH114621. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study received approval from the Institutional Review Board of all the institutions involved in the study (Fiji National University, New York Academy of Medicine and Texas A&M University).

Provenance and peer review Commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. All data are kept at the Fiji National University in FIJI, and will only be available with approval from the Fiji Ministry of Health.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Depression and other common mental disorders. Available: <https://www.who.int/publications/i/item/depression-global-health-estimates> [Accessed Jun 2021].
- The World Bank. The world bank in the Pacific islands: overview. Available: <https://www.worldbank.org/en/country/pacificislands/overview> [Accessed Jun 2021].
- Alloh FT, Regmi P, Onche I, et al. Mental health in low-and middle income countries (LMICs): going beyond the need for funding. *Health Prospect* 2018;17:12–17.
- Larkins S, Tun SSM, Kowalenko N. Strengthening mental health care in primary care in low- and middle-income countries: the Fiji experience. *J Fiji Public Health* 2018;7:35–7.
- Patel VL, Branch T, Mottur-Pilson C, et al. Public awareness about depression: the effectiveness of a patient guideline. *Int J Psychiatry Med* 2004;34:1–20.
- Hurt K, Walker RJ, Campbell JA, et al. mHealth interventions in low and middle-income countries: a systematic review. *Glob J Health Sci* 2016;8:54429.
- Ehrler F, Weinhold T, Joe J. A mobile app (BEDSide Mobility) to support nurses' tasks at the patient's bedside: usability study. *J Med Internet Res* 2018;6:e57.
- Jibb LA, Cafazzo JA, Nathan PC, et al. Development of a mHealth real-time pain selfmanagementapp for adolescents with cancer: an iterative usability testing study. *J Pediatr Oncol Nurs* 2017;34:283–94.
- Wilmer HH, Sherman LE, Chein JM. Smartphones and cognition: a review of research exploring the links between mobile technology habits and cognitive functioning. *Front Psychol* 2017;8:605.
- Kintsch W. *Comprehension: a paradigm for cognition*. New York, NY, USA: Cambridge University Press, 1998: 461.
- Kintsch W, Patel VL, Ericsson KA. The role of long-term working memory in text comprehension. *Psychologia* 1998;42:186–98.
- Patel VL, Arocha JF, Diermeier M, et al. Cognitive psychological studies of representation and use of clinical practice guidelines. *Int J Med Inform* 2001;63:147–67.
- Patel VL, Arocha JF, Diermeier M, et al. Methods of cognitive analysis to support the design and evaluation of biomedical systems: the case of clinical practice guidelines. *J Biomed Inform* 2001;34:52–66.
- Ericsson KA, Simon HA. Verbal reports as data. *Psychol Rev* 1980;87:215–51.
- Güss CD. What is going through your mind? thinking aloud as a method in cross-cultural psychology. *Front Psychol* 2018;9:1292.
- Chi MT. Quantifying qualitative analyses of verbal data: A practical guide. The journal of the learning sciences. 1997 Jul 1;6(3):271–315. "Introduction to Single Subject Designs: AB (Baseline And Intervention) Design", 2000. Available: www.msu.edu [Accessed 31 Jul 2001].
- Introduction to single subject designs: AB (baseline and intervention) design, 2000. Available: www.msu.edu [Accessed 31 Jul 2001].
- Arocha JF, Wang D, Patel VL. Identifying Reasoning strategies in medical decision making: a methodological guide. *J Biomed Inform* 2005;38:154–71.
- Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas* 1997;1:385–401.
- Osman A, Bagge CL, Gutierrez PM, et al. The suicidal behaviors Questionnaire-Revised (SBQ-R): validation with clinical and nonclinical samples. *Assessment* 2001;8:443–54.
- Opoliner A, Blacker D, Fitzmaurice G, et al. Challenges in assessing depressive symptoms in Fiji: a psychometric evaluation of the CES-D. *Int J Soc Psychiatry* 2014;60:367–76.
- Iyengar MS, Carruth TN, Florez-Arango J, et al. Informatics-based medical procedure assistance during space missions. *Hippokratia* 2008;12 Suppl 1:23–7.
- Iyengar MS, Florez-Arango JF, Garcia CA. GuideView: a system for developing structured, multimodal, multi-platform persuasive applications. *Persuasive '09 Proceedings of the 4th International Conference on Persuasive Technology*, ACM, NY, 2009.
- Florez-Arango JF, Iyengar MS, Dunn K, et al. Performance factors of mobile rich media job AIDS for community health workers. *J Am Med Inform Assoc* 2011;18:131–7.
- Gautham M, Iyengar MS, Johnson CW. Mobile phone-based clinical guidance for rural health providers in India. *Health Informatics J* 2015;21:253–66.
- Brooke J. *System usability scale (SUS): a quick-and-dirty method of system evaluation user information*. Reading, UK: Digital equipment co ltd, 1986.
- Brooke J. Sus: a retrospective. *J Usability Stud* 2013;8:29–40.
- Roberts GJ, Leckie J, Chang O. The history of mental health in Fiji. In: Minas H, Lewis M, eds. *Mental health in Asia and the Pacific, international and cultural psychology*, 2017.
- Mazaba ML, Njunju EM, Kwangu M. Suicidal ideation in Fiji: prevalence and its correlates among school-going adolescents in a global health-based survey. *Int J Public Health* 2017;9:451–6.
- WHO mental health gap action programme (mhGAP). Available: <https://www.who.int/publications/i/item/9789241596206> [Accessed Jun 2021].
- NCH software. Available: <http://www.nch.com.au> [Accessed Jun 2021].
- Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77–101.
- Creswell JW. Data analysis and representation. In: Creswell JW, ed. *Qualitative inquiry & research design: Choosing among five approaches*. Thousand Oaks, CA: Sage publications, 2013: 179–212.
- Patel VL, Groen GJ. The general and specific nature of medical expertise: A critical look. In: Ericsson KA, Smith J, eds. *Toward a general theory of expertise: prospects and limits*. New York: Cambridge University Press, 1991: 93–125.
- Iyengar MS, Chang O, Florez-Arango JF, et al. Development and usability of a mobile tool for identification of depression and suicide risk in Fiji. *Technol Health Care* 2021;29:143–53.
- Patel VL, Kaufman DR. Cognitive Science and Biomedical Informatics. In: Shortliffe EH, Cimino J, Chiang M, eds. *Biomedical informatics: computer applications in health care and biomedicine*. 5 edn, Chap 4. New York, NY: Springer-Verlag, 2021.
- Patel VL, Kaufman DR, Kannampallil TG. Diagnostic reasoning and decision making in the context of health information technology. *Rev Hum Factors Ergon* 2013;8:149–90.
- Patel VL, Branch T, Arocha JF. Errors in interpreting quantities as procedures: the case of pharmaceutical labels. *Int J Med Inform* 2002;65:193–211.